

## ACCOUNT

OFTHE


DISEASES

OF THE

ARMY.



The supplication of the state o



#### ACCOUNT

OF THE MOST COMMON

D I S E A S E S

ARMIES.

WITH THE METHOD OF CURE.

From the Original of Baron Van Swietin.

To which is added,

The Practice of Inoculation IMPARTIALLY CONSIDERED;

Popular Objections against it Consuted.

By JOHN AND REW, M.D.


WITH All

The SENTIMENTS of Dr. HUXHAM, and several of the most eminent Physicians, of both Kingdoms, relative to the PRACTICE,

As COMMUNICATED to the AUTHOR.

DUBLIN: Printed by JOHNEXSHAW.

MDCCLXVI.



The Late of the state of the st MUCCESSE.

Sugar Talanta in the observations

LIBRARY

## ACCOUNT

OFTHE

MOST COMMON

### DISEASES

INCIDENT TO

## ARMIES.

WITH THE

#### METHOD OF CURE.

Translated from the Original of

Baron V A N S W I E T I N;

Physician to their Imperial Majesties.

D U B L I N:

Brinted by JOHN EXSHAW.

## ACCOUNT

TETTOWN

MOST COMMON

## DISEASES

INCIDENT TO

## ARMIES

SINT. HITIW

### METHOD or CURE

Tabelle Official Busher'T ... V. I W. E. V. J. V. Found

Pursecián to their five entage Chartevine.

ALLE NO HON A SERVICE

Managada.



PREFACE

the find at water of the persons to be how,

office endeadour to a role bode coffice.

enits and thements.

reducidly to specify the Me. ches proper for the Core Mr the Stole, and the Diet they are to ob-



reclied the no resel of the little of

H E great and frequent Inconvenition T is cies attending a military Life, are of fuch a Nature, as make the greatest Havock, even among the most robust Constitutions. It is not therefore surprising, that an Army is seldom or ever without a great Number of Sick.

Nevertheless, it has been observed, that the Distempers mostly prevalent among the Troops are not in Number very considerable.

We have, for this reason, thought it sufficient to speak of those, whose Attacks the Soldier is most

Mag

most subject to, to describe them in such a Manner, as they may be distinguished from one another by certain Signs; and at the same Time to give an Account of such Symptoms as characterise the Decrease or Increase of each Disorder; and lastly, to specify the Medicines proper for the Cure of the Sick, and the Diet they are to observe.

It will be perceived, that, in our little Essay on this subject, it was equally necessary to be short, and to endeavour to avoid being obscure.

This Work, however, no wife regards the Phyficians, who, Masters of their Art, and in daily Practice, stand in no Need of the Assistance of these first Elements.

But it often happens, in an Army, that the Number of Sick is so great, and they are dispersed in so many different Places, that it is impossible the Physicians should go every where, and give their Attendance to each Individual. In such Cases, Necessity obliges the Sick to be entrusted to persons who cannot be expected to have the same Knowledge with those of the Profession.

For the Use of such Practitioners, not sufficiently instructed, this small Tract is designed, that, by the Signs exactly described, they may understand the Nature of the Distemper, the Conduct

they.

1-

)-

0

3.

e

8

they are to observe, and the Remedies proper to be administered.

At the End are inserted the Prescriptions reserved to in the Course of the Work. They are rendered as simple as possible, and Preserence given to those Medicines that are easiest to be procured, and easiest to be prepared.

It may not be amiss to premise some Observations, by Means of which, Sickness may in some Degree be prevented, and the Health of the Soldier preserved. We are sensible, that, in Time of War, it is not always possible to observe exactly all what we are going to say; but it cannot but be of the to know what is most advantageous, that it may be put in Practice, at least when Circumstances permit.

1. The Soldier fresh listed, and torn at once from his Family, no sooner loses Sight of his Village, but he becomes melancholy; and though a robust Husbandman, finds himself scarce able to bear the Fatigues and Inconveniencies of a military Life. It were to be wished, that he could be used little by little, to this new Kind of Life; but in the mean Time nothing is better, than to procure him all Kinds of Amusement and Diversion.

2. Garden

to character selection themself all the property

- 2. Garden Stuff and fresh Greens are a wholefome Nourishment for the Soldier; ripe Fruit are
  equally good, and never hurt but by their Abuse;
  but unripe and acrid Fruits are very hurtful.
  The Use of Garden Stuff and Fruit prevents the
  Scurvy, and even cures those already attacked
  with it.
- 3. The Choice of Water is effential. If there is none to be found absolutely pure, the Preference is to be given to that least loaded with heterogeneous Parts. There is an easy Way to distinguish Water that is pure from that which is less so, by Means of oleum tartari per deliquian. If you let sall a few Drops of this in a Glass of Water, if not pure, it becomes instantly thick, and milky, whilst the same Operation produces only a small Cloud in that which is more pure.

River Water should not be drawn near the Banks; that which is in the Middle of the Stream is always best.

Sometimes indeed Troops are reduced to the fad Necessity of drinking bad Water; in which Case it may be much corrected, by mixing a certain Quantity of Vinegar. Six Ounces, mixed with three Quarts of Water, will render the Drink even more agreeable.

n

a

le-

re

e;

hie

ed

re

e

-

h

y

t

1

The Water may also be rendered less hurtful by steeping in it some Pieces of the Root of the calamus aromaticus. This Root is found every where, especially in marshy Places, where commonly the Water is worth.

4. The Soldier ought to be well cloathed and covered; his Shoes of a thick and ftrong Leather, and the Thread they are sewed with well covered with Wax: If all the Seams could be well waxed, the Water would be prevented from penetrating.

as much as possible. That which appears such is sometimes quite the reverse, because the Waters are but at a little Distance below the Surface. It is, however, very easy to know that, by digging the Ground, or even merely by examining the Wells of the neighbouring Villages.

If the Water is high in the Wells, the Soil is damp; if the Water is low, the Soil is dry.

It would likewise be proper to avoid the Neighbourhood of thick Forests; they prevent the Wind from penetrating, and render the Air of the circumjacent Parts damp and close.

But if there is a Necessity of encamping in a damp Ground, the Soldiers must have their Straw changed changed oftener than usual. The Officers will be much benefited by spreading a waxed Cloth under their Bed.

In Times of Rain, the tighter the Tents are extended, the less it penetrates: Small Trenches dug round the Tents contribute to the Dryness of the Spot where the Soldier lies, because they receive the Water that falls from above.

6. When an Army remains long in the same Camp, the unwholesome Effluvia from so many Bodies always occasion Sickness, unless there happens to be violent and frequent Winds. These Exhalations are mostly to be seared in a hot and moist Air. The changes of Camps, on this Account, contribute much to the Health of the Soldier, more especially when the Bloody Flux prevails. Hence arises one Reason the more for avoiding the Neighbourhood of thick Forests, which prevent the free Passage of the Winds.

7. Nothing is more prejudicial to the Soldier, when heated with Work, than to strip, expose himself to the cool Air, and greedily drink cold Water, and especially Well-water, which commonly is very much so. River Water is less hurtful, as the Rays of the Sun, to which it is contantly exposed, prevent its being so cold.



OF THE MA neit , orer

To Man relatives of the control

# DISEASES

of THE signed Affacts

### ARMY.



reallised, they four ince they

F the Troops encamp in the Spring, especially early in the Season, there will infallibly be many fick. The Difeases that principally prevail at that Time are, very troublesome Coughs, fore Throats, Pleurisies, Peripneumonies, and Rhenmatisms.

These Distempers, though not contagious, do not admit of much Motion in the Patients, wherefore it should be endeavoured to have the Hospirals nigh Hand, and if the State of the diffemper requires it, to bleed the Patient, before he be moved, the Delay of which may produce very troubletome Confequences,

Sometimes affo Intermitting Fevers prevail during this Season, but they generally are less obstinate, than those that happen in Autumn. In Spring they are commonly Tertians, or Quotidians; but seldom Quartans, unless in those subjects, who have been before attacked during the Winter, and in whom, properly speaking, they are only Relapses.

## COUGHS.

OUGHS are more troublesome than dangerous: But if they continue any Length of Time, and are neglected, they sometimes degenerate into a Philis Pulmonalis.

for his common Drink, and drink it Luke-warm; it will be of Service to add a Fourth Part of new Milk.

the Seatons affere

The Patient must abstain from Wine, from salt and acid Food. Rice and Barley Broth, and new Milk with the Yolk of an Egg, will be sufficient Nourishment.

If the Cough becomes too violent and troublefome, fo as to prevent the Patient from Beeping. he may take the Prescription No 2.

If the Cough is attended with a Feyer, Bleeding will be necessary to prevent an Inflammation. which is then to be apprehended.

When the Cough diminishes, and the Excretions, which before were without Confistence, become thick, and are easily expectorated, the Diftemper is at an End.

#### the raid the their constitution as the factories SORE THROAT. a Sher or twee the Diferio active & the Great of

some of the tention provided freshold to the one

individual transfer fregisties and research com F the Action either of Swallowing or Respiration fuffer an Impediment, attended with fensible Pain, and the Cause is in the Neck, or Throat, the Disorder is called a Sore Throat, (Angina.)

in even with a property of the state of

It is a very dangerous Distemper; and sometimes even mortal; it is known to be fo, when respiration is greatly impedited, the Voice extremely sharpened, and the Anxiety very confiderable. A large Bleeding is immediately requifite, and Cupping Glasses are to be applied, around and on the Nape of the Neck, which often give instant Relief: The Drink No 1, must be:

f

kept constantly in the Mouth warm; and the poultice No 3, applied and continued hot on the Neck, Night and Day.

If the Patient can swallow, let him take every Hour a Cupsul of the Drink No 1. made hot, with the Addition of twenty Grains of purified Nitre to each Pint. If the Redness appears on the Neck or Breast, the Patient often recovers. Otherwise this Disease is frequently mortal; but it is not common. The following Species is much more frequent.

One of the tonfils grows red, swelled and painful, and the Pain commonly extends itself to the internal Part of the Ear of the diseased Side. In a Day or two, the Disease attacks the Gland of the other Side, while the Swelling disappears from that which was first affected. Sometimes the Pulse is hard and quickened, and sometimes not at all.

In the first Case, the Urine appears of a deeper Colour than it is in Health; Bleeding is then necessary, and if the Redness, swelling of the Throat, and Difficulty of Breathing do not give Way, it will be proper to repeat it.

In the fecond Case, viz. if the Pulse is natural, there is no Necessity for Bleeding, unless the Patient be of a replete Habit.

The Nourishment must be restrained to light Broth, to which may be added Rice or Barley! Cream.

Let the Patient take a Tearcup of the Drink.

No 4. every Hour warm, unless he sleeps: and let him often keep in his Mouth some of the medicated Tea, No 5. warm, with which he is also to gargle his Throat.

Next Day let him be purged with the Draught No 6. which, if the Disorder does not abate, he may take it again after two Days, continuing, in the mean Time, the Use of No 4. and No 5. till he can swallow freely, and the Redness hath disappeared from the Iaside of the Throat.

If the Disorder has continued too long, before Relief was applied, or the Inflammation is too confiderable, (to give Way) Suppuration will enfue.

That there will be a Suppuration is known, by the Swelling and Redness in the Throat continuing above three Days, without any Abatement. In this Case, the Patient is to keep continually in his Mouth some of the Decoction No. 7. warm, which likewise may be lightly injected into his Throat.

The

The Catapialin No 8, is to be applied, and kept Night and Day round about his Throat.

If the Swelling of the Throat then begins to grow fort, the Abscess will soon burst: If there appear a little spot elevated and white, the concealed lancet (Pharyngotomus) may with all Safety be used, to give by this Means a free little to the Pus.

The Abcess being either broke of itself, or opened by the Lancet, the Gargarism, No 9, must be frequently used, and the cure will soon be compleated.

When it happens that the Swelling prevents Deglutition entirely, a Glyster of twelve Ounces of new Milk, and fix Ounces of Barley Water is to be administered every four Hours, which the Patient must retain as long as he possibly can. By this Means he may be sustained till the Abscess breaks.

There is still another Species of Sore Throat, which, in the Beginning, is easily cured, but by Neglect, degenerates into a Sort of Mortification, and corrodes all the Parts attacked, causing at the same Time a most horrible Stench.

terrogical administrational associations

There

There appears on the Tonfils, on the Palate, on both Sides of the interior Part of the Mouth, or within the Lips, one or feveral white Spots, fometimes yellowifh, and even brown, according as the Diforder is more or less violent. The Places round these Spots are much inflamed and painful. It sometimes happens that the patient has no Fever; and the Swelling is never to confidenable in this Case, as in the before described fore Throat.

The last Species generally goes off pretty foon, by rubbing every two Hours lightly the Parts affected with a Pencil dipped in the Mixture N° 20, and using for a Gargarism a simple Insusion of Elder Flowers. It is proper the Patient should drink some Cupe of the same Insusion about sour Times a Day.

3

:3

is

B.

ti,

re

It is to be observed that the said Spots increase very suddenly, when the Stench of the Mouth is great: In that Case, the Quantity of sp. fal. mar. must be augmented, to stop the Progress of this Evil.

1953 I will be received with extent on the

distributed in the same of the same

Cit Ligital theory is a west blood to the

The

## The PLEURISY.

or estate the hins the conservation while Spore

The State of the S

THE PLEURISY is known by a sherp Pain, and Stiches felt in the Breast and attended with a Fever.

This Pain is increased by Inspiration, and leffened by Expiration; and by holding the Breath; the Pulse is generally hard, as in all acute and inflammatory Diseases: In violent Pleurisies, the Pain is sometimes so sharp, that the Patient scarce dares to draw his Breath, the Face grows livid, and hes feels himself just suffocating; in this State the Pulse is small and low.

by the violent Pain: in some Cases, this Cough is dry, without any Excretion; in others it is, from the Beginning, attended with Excretions; less Danger is to be apprehended in the last Case than in the first.

no we blook is list shi become the leaves them

Though the Sides of the Breast are most commonly attacked in this Distemper, yet the anterior and posterior Parts may be equally affected.

If the Pain is most sensibly felt externally, and chiefly upon touching the Part, the Disorder is commonly named a false Pleurisy, and is to be treated as follows.

Bleeding

Bleeding is the first and chief Remedy: let 12 Ounces of Blood, or even more, if the Subject is replete and robust, be taken from the Arm on the Side where the Pain is. While the Vein is open, let the Patient respire strongly and cough. Bleeding commonly lessens the Pain, and sometimes carries it off altogether.

Some Hours after the Bleeding, the Glyster, No 11. ought to be given.

Flannels dipt in the Fomentation No 12. are to be constantly applied to the Place where the Pain is felt; and as it would be troublesome to apply this Topic during the Night, a Plaister of Labdanum spread on Leather or Linen, may be applied in its Room.

In the Morning this Plaister must be taken off, the Part rubbed with the ung. alther and the Medicine No 12. Let the Patient take every Half Hour, if he be awake, a Spoonful of the Mixture No 13. and drink after it a warm Cupfull of the Decoction No 1. to each Pint of which one Ounce of Honey should be added.

It often happens, that the Pain, which after Bleeding was much leffened, or quite ceafed, returns with as much Violence as at first: In this Case a second Bleeding is necessary, but commonly not so large as the first, which always ought to be very copious: If after this the Pain still returns very fensibly, you must bleed a Third Time, and sometimes even a fourth, according to the Violence of the Distemper.

It must however be observed, that slight remains of Pain, and such as but little impede Respiration, do not require fresh Bleedings: the Patient might by them be too much weakened, and rendered languid a long Time after.

Bleeding therefore must only be repeated in those Cases where the Pain is violent enough to obstruct the Breath considerably; the Velocity of the Pulse increased commonly at the same Time shews the Necessity for this Evacuation.

It is a good Sign, when the Pain changes its Place and affects the Clavicles, the Shoulder-blades, the Shoulders and Back; and that this new Pain requires no Bleeding. These Changes generally happen about the fixth Day: It is then sufficient to chaff the Parts in Pain lightly, and then anoint them with the ung. alth:

Let the Nourishment be light, and consist of thin Broth, roasted Apples, and well fermented Bread: The Decoction No 1. or plain Barley Water with one Fourth of new Milk, may serve for common Drink. If the Belly be constipated, the Glyster No 11. may be repeated. As foon as Respiration grows easier, and the Pain is considerably diminished, it will be sufficient to give every two Hours a Spoonful of No 13, with a Cup-sull of the Decoction No 1. warm after it.

But if, notwithstanding several Bleedings, the Pain continues, without any sensible Diminution; and above all, if the Rattling of the Breast, and Want of Expectoration shew the lungs to be filling, a Blister is to be applied to each Calf of the Leg.

A sharp Blister applied on the painful Part has often been attended with very good Effects, even when repeated Bleedings had procured no Cestation of the Pain of the Side.

Care must be taken, in this as well as in all instantant matter Distempers, that the Patient be not kept too hot, and that the Air be often renewed.

When the Illness begins to grow milder, by the Use of the above Medicines, fresh Symptoms will appear, which shew the morbific Matter is concorded and ready to be expelled.

Great Care must then be taken, not to present its Course, but on the contrary to facilitate it by all the Means that Art can suggest,

Bleeding Piles are of Service; a white, rediff, or sometimes a brownish Sediment in the Urine is a favourable Symptom: To encourage its Continuance, the Sick must drink plentifully.

If in the Progress of the Distemper, after the Symptoms grow milder, the Excrements are yellow, and bilious, it is a good Sign; but at the Beginning it affords but a bad Prognoftick.

In general this Disease is carried off by Expectoration, especially if the Excretions are plentiful and diminish the pleuritick Pain, and above all, if they are ripe and resemble Pus. Sometimes they are glewy, tenacious, and sanguineous; but there is nothing to fear from them, if the Pain abates, the Fever diminishes, and Respiration becomes more free. Be cautious however not to repeat the Bleeding, which, under these Circumstances, must be prejudicial. Sometimes the Excretions are yellowish, and mixed with Streaks of Blood, and this also is a favourable Sign.

Laftly, it may be laid down as a general Rule, That Expectoration is to be eftermed a happy Symptom; whenever the Excretions are thrown off with Ease, they diminish the Pain and Fever, and render the Respiration easier:

In such a State of the Expectoration, the Mixture No 13, must be laid aside, and the Linctus No 14 substituted in its Place; two Spoonfuls of which are to be given once in an Hour, and leifurely swallowed, with a Cup of the Decoction No 1, warm after it.

If the Expectoration does not continue, but suddenly stops, and there comes on a Rattling in the Breast, attended with Anxiety, the Danger is very great; Blisters on the Legs must be applied immediately, the Powder No 15 be given every four Hours, and the warm Decoction No 1 sweetned with a little Honey, be drank plentifully, till the Spitting comes on again, and the Breast is relieved.

There comes on sometimes, but not often, a painful Swelling behind the Ears, or on the Thighs, which Pain is followed by a Diminution of that of the Breast: In this Case, the immediate Use of the Cataplasm No 8. or such like, is necessary, to ripen the Swelling, which is to be opened with the Lancet, as soon as it comes

to a Head and be dreffed afterwards as an Ulcer.

The Violence of the Distemper may be fo great, that the most efficacious Remedies are insufficient to overcome it, and to expel the morbific Matter. In this Case, Suppuration, which is always dangerous, comes on, and most commonly the Illness degenerates into a Consumption, unless Means be found to evcuate the already formed Pus.

That this is the State of the Case, may be known by the following Symptoms.

The Pain is stubborn, and yet less violent than at the Beginning: It is attended with a dry Cough, or with unripe Excretions; there is a continual Quickness of the Pulse, which increases towards Night, or whenever the Patient takes any Nourishment; the Cheeks and Lips become red; he has frequent Shiverings and Night-Sweats; the Urine is frothy and pale, and he foon becomes extremely weak and lean. The Abcess, formed in the Lungs, is fometimes evacuted by Excretions: When they begin to appear, and are purulent, the Infusion No 16. sweetened with a little Honey, is to be given every Hour: Broths, in which fresh Chervil, lettice, and Parsley Roots, are boiled, may ferve for Nourishment; and for Drink, Barley-Water, with a Fourth Part of new Milk.

Milk, both to be continued till the purulent Matter is quite evacuated.

But this does not always happen; the Pusis often formed in a Bag; and in that Cafe it must be attempted to draw out this Collection of Matter.

t,

nt

r.

n-Is

oe

\*4.

VR

an

h,

al

ab

u-

he

he

es

ed ti-

u-

tle

in

ts,

ew

lk.

A small Plaister applied, and made to stick closely to the most painful Place, will be very proper in the Beginning of the Illness; because if the Pleurify should degenerate into an Abscess, the Collection of Matter will point towards that Place:

When an Abcess is known to be formed, by the Signs that have been described, the marked Place is to be corroded by a light Claustic; and when open, Care must be taken to keep up the Supparation. In such a Case, there is Reason to hope; as the Resistance is the least at this Place, that the Matter collected will take its Course, and be discharged by it. For these Collections are often lodged between the Pleura and the adjacent Parts.

For the same Reason, a Seton on the Spot is successfully used; and the Pus has often been feen to discharge through such a Passage procured by Art.

If the Matter contained in the Abscess cannot be drawn into the external Parts, it will occasion a C 2 swelling

fwelling of the Pleura towards the Cavity of the Breaft; whereby the Lungs will be oppreffed, the Anxiety daily increase, the Pleura burft, all the Symptoms fuddenly disappear, but come on again foon, and the Pus fall into the Cavity of the The second of the second of the second Breaft.

Under these Circumstances, no other Method can be used but the Paracenthesis, to discharge the Breaft from the Pus there lodged, and prevent a mortal Confumption.

During this last Trial, the Use of the Medicine No 16. is to be continued. If, during the Course of the Distemper, the Patient can get no Sleep, a Pint of Emulsion No 17. is to be administered, to which may occasionally be added an Ounce of the Syrup of white Poppies, or more if neceffary. make the first the method was in the first of the second

ME TO SHOW ME THE STATE OF THE asuassin envision as character less of bourse soul mosaide of the distribution of the adjacent fund

Puchasio I'm st 22 to manger from it washing

PERIPods or other ways

### PERIPNEUMONY.

he

he he in

he

N. 17

od

he

a

di-

he

no

if4

an

if

100

T HIS Diftemper is, properly speaking, and Inflammation of the Lungs: It is dangerous, and even more so than the Pleurisy itself, which sometimes degenerates into a Peripneumony, when the Patient is forced by Excess of Painto keep in his Breath.

A Difficulty of Breathing, the Load and Oppression of the Breast, and an acute and continual Fever, indicate a Peripneumony. In this Distemper the Patient seels no Pain; or if he does complain of any, it is of such a dull one as distinguishes the Peripneumony from the Pleurisy; which last, in Inspiration, gives the Patient a very acute Pain.

The Pulse is not so hard in the Disease we are now describing, as in the Pleurisy and other inflammatory Distempers, but, on the contrary, is generally found much softer.

If the Peripneumony is violent, there immediately comes on a great Weakness, the Pulse becomes small, soft, unequal; the Respiration is short, frequent, difficult, and accompanied with a continual Cough; the Patient cannot lie down for Fear of Suffocation, but is obliged to sit upright

C 3

in his Bed; his Face, Eyes, Tongue, and Lips, become red and inflamed: These Symptoms are followed by an insupportable Anxiety, and soon after by a Dilirium, and Death.

All the Signs, therefore, that we have recited, give us a very bad Prognostick.

A greater Hardness in the Pulse, a less Difficulty in Breathing, more Ease in lying down, less Redness and Swelling in the Face, the Eyes, and the Lips, are, on the contrary, favourable Symptoms.

This Distemper requires immediate Help, for very soon it brings the Patient into evident Danger.

You must begin by a large Bleeding in the Arm, and repeat it in the same Manner as in a Pleurisy, if the Anxiety and Difficulty of Breathing do not diminish. If the Blood, when drawn, remains studid and thin, and scarce coagulates at all, and if after the Bleeding the Respiration is not freer, it is a bad Sign, which indicates, that the thicker Parts are retained in the Lungs, and the thinner only discharged. In this Case a fresh Bleeding would produce no Effect, and only evacuate that Part of the Blood which is least thick, and could still have made its Way through the Lungs.

ps,

are

noc

ed,

Dif-

wn,

res,

ble

for

an-

rm,

ify,

not flu-

d if

eer,

ker

ling

hat

uld

ome

Some Hours after the Bleeding, it will be proper to give the Glyster No 11. Fomentations, Ointments, and Plaisters, may be applied to the Breast, but it must not be expected that these Sort of Things will be attended with as good Success as in the Pleurify.

It will be better to apply frequently to the Patient's Mouth and Nostrils, a Linen or Sponge, foaked in warm Water, the Vapours of which may, together with the Air, enter by Inspiration into the Lungs.

The Diet ought, as in the Pleurify, to be extremely light, and the Broths fill thinner.

Let the common Drink be the Decoction Nor, or Barley-Water; but instead of mixing Milk with it, add to each Pint Half an Ounce of pure Honey.

While the Patient is awake, let him take every Half Hour a Spoonful of No 13. and drink after it a warm Cup of the Decoction No 1.

If, on the Use of these Medicines, the Anxiety diminishes, the Respiration becomes freer, the Fever less violent, the Pulse more vigorous and equal, the Tongue moist, and every Part of the Body, even to the Extremities, of an equal Heat; if, above

bove all, the Skin is moist and soft, we may hope for the best, and nothing more is required than a Continuance of the same Means, as the Instammation of the Lungs is beginning to be resolved, and gradually to decline.

But Things seldom take this Turn, unless the Distemper is not violent, the solid Parts are supple, and Relief has been applied from the Beginning. It oftener happens that the Matter of this Disease, is evacuated by Expectoration.

The Spitting must therefore be carefully attended to; and it is a very bad Sign, when at the same Time the Patient does not expectorate at all, and has a Difficulty to breathe, with a Rattling in his Throat. The Spittings are good, if discharged speedily, copiously, and easily. They should be of a proper Consistence; sometimes they appear yellow, and streaked with a little Blood; which ought to give no Uneasiness, for Excretions of this Sort are always good, and grow white in Time.

The Effect they produce is remarkable, by the Diminution of Anxiety, the Freedom of Respiration, and the Alteration in the Pulse, which grows stronger and fuller.

P

Let the Patient, at that Time, take two Teafpoonfuls of the Linctus No 14. and after swallowing them softly, drink a Cup of the warm Decoction No 1.

Nothing more is to be done under these Circumstances, and Bleeding, Purging, or exciting Sweats, would be detrimental.

We ought chiefly to guard against the cold Air, and cold Drinks, for either the one or the other will stop the Expectoration, and thereby throw she Patient into imminent Danger.

If a Suppression of the Spitting should happen, and the Anxiety be followed by the Rattling of the Throat, Blisters are to be applied to the Legs, the Powder N° 15. is to be given every four Hours, and the same Decoction N° 1. to be used as ordered for the Pleurisy. The Patient is also to inspire by the Mouth and Nostrils the Vapour of warm Water.

It sometimes happens, that, during the Course of the Distemper, the Patient voids by Stool, a yellow and bilious Matter, and is relieved by it. This is also a favourable Sign, as has been observed, in speaking of the Pleurisy.

A large and thick Sediment in the Urine, at first red, afterwards turning white, is likewise a good Symptom. When this happens, the Patient must drink plentifully as in the Pleurisy. Nevertheless, it rarely happens that the Cause of the Evil is discharged merely by Urine; the Spitting which generally comes on about the same Time; contributes greatly to the entire Cure.

When the Patient, by Means of these Evacuations, begins to find his Breast disengaged, he may have his Broth somewhat stronger; but he should constantly take but little at a Time and often, that the Lungs may not asresh be overcharged by a Chyle, both too crude and too copious. Sometimes a plentiful Bleeding of the Nose gives Relief to the Patient; but this seldom happens.

If none of the Evacuations here described are observed within the Space of a Fortnight, if the Fever continues pretty strong, and the Coughdry; if the Heat extends to the Extremities of the Body; if the Pulse is quick, soft, and wavering; if a Difficulty of Breathing and Shiverings accompany these Symptoms; if the Cheeks and Lips are red, the Thirst great, and lastly the Fever stronger towards Night, it is certain that the Instammation is turning to an Abscess.

The

fe

li

ti

tl

V

B

ti

P

LE

a

ıt

-

e.

g

3.

y

d

18

a. -

f

e h

f

-

5

d

•

e

The Indications of an Abscess already formed in the Lungs, besides the Symptoms already described, are as follows: A dry continual obstinate Cough, which increases when the Sick moves, or takes any Nourishment; he can only lie on the affected Side, without its being possible for him to lie on the other; he has periodically a little continued Fever, which augments whenever he eats, drinks, or stirs, and is attended with a Redness of the Lips and Cheeks; he has no Appetite, but a violent Thirst, and Complaints of Night-Sweats, especially of the Head, and she Upper Part of the Breast; the Urine is spumous; he is greatly emaciated, and extremely weak.

While the Abscess remains whole, the purulent Tumour increases more and more; it presses upon those Parts of the Lungs which as yet are found; it obstructs the Respiration, and after the most terrible Anxieties, suffocates the Patient.

It is therefore effential, that the Abscess should break and be maturated, in order that the Pus be evacuated. But it may happen to break in such a Manner as to disperse the Matter into the Breast, and occasion an Empyema, which almost always proves mortal. This is known to be the Case, by the sudden Diminution of all the Symptoms, sometimes attended with slight Faintings, and the total Suppression of purnlent Spittings. It is because the

the Abscess is in effect burst, that the Symptoms proceeding from the Distention of the purulent Bag immediately cease; but the Matter spread in the Cavity of the Breast, growing every day more copious, and more acrid, soon occasions new Symptoms, worse than the preceding ones.

The Paracenthesis is the only Means of Help left: But as the Ulcer has already corroded the Substance of the Lungs, the Success will be very doubtful: And even though the Pus is evacuated, the sick Person generally dies after the Operation.

Much more is it to be wished, that the Abscess may break in such a Manner, as to let the Pus fall into the Bronchi or Air-Vessels of the Lungs, that so it may be evacuated by Spitting.

When this happens, it is to be feared, left the Bronchi be totally filled and stopped by the Quantity of Pus spreading itself instantly, and all at once, and occasioning thereby a Suffocation: But if the Pus that falls in these Vessels can still be discharged, the Patient often recovers, though the purulent Consumption is always to be apprehended.

The following are the principal Succours which Art has found out to forward the opening of the Abscess in the Bronchi, and the Evacuation of the Pus by Expectoration.

As

ċ

T

22

m

pe

fc

21

dit

pe Ex

to

ns

nt

n

re

0-

p

10

y

t-

e-

ſŝ

II

at

ne ·

n+

at

ut

b

**C**-

h

ne

13

As foon as the Symptoms of an Abscess, as described above, are perceived, let the Sick inspire continually by the Mouth and Nostrils the Vapour of warm Water, to soften and relax the Parts.

Let him take fatter Broth, and in larger Quantity, than before, that the Stomach being filled, the Descent of the Diaphragm may be more difficult, and the Abscess more compressed. The Patient must be excited to cough, by applying warm Vinegar to his Nostrils, or to cry with a loud Voice. This may procure a Chance of breaking the Abscess, which, if the Strength of the fick Person can bear it, may be still forwarded by giving him an Airing in a Carriage, in a rough Road, where he may be well shook.

As it is impossible to know exactly the Moment when the Abscess will break, you must repeat from Time to Time the Attempts here described.

If, when the Abscess is broke, the Excretions are purulent, white, and smooth; if the Fever disappears, or diminishes considerably; if the Appetite returns, the Thirst ceases, and lastly, the Excrements are solid and natural, there is Reason to hope the Sick will recover.

If, on the contrary, the Excretions are stained of different Colours, with a bad Smell; if the Fever doth not cease, or having ceased, returns again; if the Thirst remains, and the Appetite doth not increase, it is to be feared the Patient will sink under it.

When the Abscess of the Lungs discharges itfelf by Means of purulent Excretions, a little Rice, or Oats boiled in Milk, affords an excellent Nourishment; but Care must be taken, that the Sick take not too much at once, but little and often.

The Infusion N° 15, with a Third Part Milk and a little Honey, will be a proper Drink. He is to take thrice a-Day the Powder N° 18, and as the Lungs have been fatigued by a continual Cough during the Day, some Relaxation ought to be procured in the Night; wherefore let the Patient take two Pills N° 19.

If he is fomewhat costive it is not amis, but if he remains so seweral Days, the Glyster No 11. is to be given. If the Excretions diminish little by little, Appetite comes on, Strength increases, and the Patient gets rid of his Fever, a quick Cure may be expected.

When

ed

no

fm

it

en

ftr

to

to

to

ced

the

Ac

Ad

tio

Pa by

fre

hir

th

eo.

de

When the Excretions are confiderably leffened, the Powder N° 18. and Infusion N° 16. are no more to be used; instead of which, three small Spoonfuls of the Linctus N° 20. and after it three Cups of the Insuson N° 21. may be taken thrice a-Day.

ed

he

rns

ite

ent

it-

tle

elhat

ba

lilk He

25

ual

the

t if

is

by

nd

ire

en

If, notwithstanding this, the Cough returns stronger towards Night, the Patient may continue to take the Pills No 19, which otherwise are also to be left off.

If, after the Abscess in the Lungs hath begun to be evacuated by Excretion, this Evacuation should suddenly stop, an extreme Anxiety succeds, together with a Rattling in the Breast, and the Sick is in very imminent Danger. This Accident is commonly caused by the imprudent Admission of cold Air, or by some violent Emotion of the Mind, as Anger, Fear, or such like.

A speedy Relief must be given, by causing the Patient to inspire the Vapour of warm Water, both by the Mouth and Nostrils, making him drink freely of the warm Insusion No 16. and giving him every four Hours the Powder No 13. until the Breast be disengaged, and Expectoration comes on again; after which the Use of the Powder is to be discontinued.

It will be of Service to apply Blisters to the Calves of the Legs, as has been recommended in the Pleurify.

When the purulent Matter is reforbed by the Veins, it sometimes occasions a sudden deposit of Matter in other Parts of the Body, and there causes an Abscess, viz. about the Ears, the Arms, or Thighs. The Load of the Breast ceases at the same Time; and the same Remedies are to be used, and the same Rules observed, as in the Case of an Abscess after a Pleurisy.

As the Inflammations of the external Parts may degenerate into schirrous Hardnesses, the same Accident is to be feared in Inflammations of the internal Parts: For after a Peripneumony, there fometimes remains in the Lungs a schirrous and callous Hardness, in which Case there is almost always an Adhesion to the Pleura. Respiration continues in that Case difficult for the Rest of Life, attended with a little Cough, chiefly after Meals, and after Exercise; and there is no Appearance of any of the Indications of an Abscess we have been describing. This Accident can rarely be got the better of; and the little Relief that may be afforded is scarce to be expected in a military Life, unless it be for the Cavalry, by riding on Horseback.

ar

th

in

u E

ы

Laftly, if the Peripneumony be so violent, that the Remedies have no Effect, Gangrene and Death are the Consequences. This is foreseen, when the Patient labours with intolerable Anxiety, falling into extreme and fudden Faintnesses, with an unequal, feeble, and very quick Pulse, and the Excretions without Confiftence stinking and black. All these Signs indicate a speedy and inevitable Death.

多。在1980年,1980年,1980年中华的1980年,1980年以

The second of th

con a longer of the best like the mile

the Day of State of the State o 

state of the parties of the state of the sta THE HOLD THE STATE OF THE STATE

the second of the last of the second of the

The second secon

to proceed the supplied the problem of the problem

and the second s the state of the s

and the second of the second o

3

e

f

T e

e

e.

e

е d

n

of

1: )-

S

-

it i-

g

y .

to a first the second state of the last of D3 RHEUMA-

ENTRE PROPERTY OF THE PARTY OF

BEN BUILD

# RHEUMATISM, AND RHEUMATIC PAINS.

THE general Cause of the Rheumatism is the sudden Exposition of the Body to the Cold, after being considerably heated, either by Work or the Weather; especially if a Person being overcome by the Heat, throws off his Cloaths, and rests himself in a damp and cold Place.

The Soldier is most frequently liable to this Disorder, when heated by Fatigue, and exposed to Rain, he is obliged to wear his wet Cloaths, without changing them.

The cold Nights, which in Spring and Autumn fuceeed very warm Days, are also a Cause of the Rheumatism. This Distemper begins with an universal shivering, followed by Heat, Thirst, Uneasiness, and Fever. After a Day or two, and even sometimes sooner, the Patient seels a sharp Pain, not confined to one Place, but moving from one Limb to another, viz. at the Wrists, the Shoulders, the Knees; and so different Parts of the Body are in this Manner successively affected; and the Articulations, that are attacked, become red and swelled.

Sometimes, in this Disorder, the tendinous Expansions covering the Muscles are attacked, and occasion an excessive Pain on the least Motion of the Part affected. Sometimes the Fever ceases in a sew Days, though the Pain continues. In general, it is a very troublesome Disorder, especially when it attacks the Loins, as the Sick is then confined to a supine Posture, without Motion, and, as it were, like a Log. It often passes from the Loins to the Hips, or upper Joints of the Thighs; where if it remains long fixed, the Cure becomes very difficult.

When the Pain suddenly and often changes from Place to Place, it is to be feared that the Cause of the Illness is drove inwardly, and attacks the Lungs, or Brain, which would be attended with the greatest Danger. This Accident is discovered by a Delirium, or a violent Oppression of the Breast succeeding a Cessation of Pain in the Extremities.

This Disease is seldom mortal; but the Violence of the Pains, and their Continuation upon an improper Treatment, induce us to employ quick and efficacious Remedies. When the Disorder is neglected, it often happens, that the Articulations are deprived of Motion, and there remains for Life an incurable Stiffness of the Joint (Anshylosis.)

The

The Method of Treatment is as follows.

Take ten Ounces of Blood from the Arm of the

Let Flannels imbibed in warm Fomentation No 12. be constantly applied to the Part in Pain.

The Diet ought to be light, confisting of small Broth, with Decoctions of Barley, Oats, or Rice, and with roasted Apples.

For common Drink, use the Decoction No 1. or Barley-Water with a fourth Part Milk.

Two Spoonfuls of the Mixture N° 22. with a Cupful of the Infusion N° 23. warm, may be given every Hour, unless the Patient be asseep.

The following Day let him have the Glyster No 11. and constantly continue the Medicines No 22. and 23.

If the Pain does not give Way, and the Fever continues, the Bleeding is to be repeated the next Day, the Fomentation, No 12, and the Medicines No 22, and 23, are to be continued; after which, the following Morning, let the Patient take the Purge No 6, omitting during this Day the Use of

N° 22. and 23. and taking at Night the anodyne Draught N° 24. Let him afterwards continue two Days longer the Use of N° 22. and 23. and on the third Day repeat the Purge N° 6. and at night the Draught N° 24.

By this Method, we commonly get the better of this Diforder. If a good Deal of a Brick-coloured Sediment appears in the Urine, attended with a gentle breathing Sweat, it is a good Sign.

It will then be sufficient to compleat the Cure, to keep the Patient warm in Bed, and make him take the Decoction No 23.

But if, after the Use of these Remedies, the Pain still continues, and the Part affected grows red, Leeches are to be applied upon it.

Sometimes the Fever ceases, the Patient appears recovered, but the Pain still roves from one Joint to another: In this Case let the Patient take Half a Drachm of Venice Soap made into Pills, Morning, Noon, and Night, drinking after it as Ounces of the Insusion No 2.3. made warm. He must be kept from the Cold, and the Articulations lightly rubbed with a Piece of dry Flannel.

It happens likewise sometimes, that the Patient, tho' otherwise recovered, has a fixed Pain left about the Articulation of the Hip. Apply in that Case a Blister about the Size of a Crown Piece

upon the Part for twelve Hours, then take it off, and Pierce the Bladder it has raifed that the lymph amassed may flow out, and cure the Wound by Means of the Plaister, called empl. allum cocum.

Eight Days after the Place where the Blifter was applied is healed, apply another, and proceed as before, and if the Pain is then not entirely vanished, this Operation may be repeated four Times. Observe, when you take off the Blifter, only to pierce the Vesicle, and not take off the Epidermis, for the Place thus made bare, would be extremely painful, and without any advance towards the Cure.

Those who have suffered of this Disorder in Autumn, must take Care during the following Winter not to expose themselves to the Cold, and the Injuries of the Season, for they would certainly have a Relapse.

When by the Pain continuing long fixed in the same Place, the affected Articulation begins to shiffen, let the Part be twice a day held over the Steam of hot Water, then well wiped with hot Linen, lightly rubbed, and anointed with ung. alther.

with the party the manufacture

FOR THE PROPERTY AND TH

VIGO Q I off

Course to Kill religion

## INTERMITTING FEVERS.

A FEVER is known by a Quickness of the Pulse, usually attended with Lassitude, Languor, Weakness, Thirst, and several other Symptoms.

When after a Fit of several Hours, it sensibly diminishes with all its Symptoms, and at last absolutely ceases, but in such a Manner as to return again, it is called an Intermitting Fever.

This Fever has different Appellations according to the Length of the Interval between the Fits.

If it returns every Day, it is called a Quotidian; if there is a Day between each fit, it is called a Tertian; if the Return is after two Days free, it is named a Quartan.

The Intermitting Fever comes on with gaping, Lassitude, Debility, Colds, Shiverings, tremblings, Paleness of the Extremities, Anxieties, Sickness, and sometimes Vomiting. The Pulse is feeble and small, and the Thirst pretty great. Heat succeeds after some Time; it insensibly augments till it becomes extreme. The Body then grows red, the Anxiety diminishes, the Pulse is fuller

fuller and stronger, the Thirst excessive, and the Patient complains of a violent Head - Ach, and Pain in all his Limbs. Lastly a general Sweat succeeds; all the described Symptoms diminish, and the Patient often falls into a Sleep, after which wakes without Fever, his Pulse is natural, and there remains nothing but Lassitude and Weakness. Sometimes during the hot sit of the Fever bilious Matters are thrown up with a Perception of Relief.

The Urine after the Fever, or during the Sweat, is reddish and frothy; and as soon as it is cold, there appears on the Top a Pellicle, adhering to the Sides of the Vessel, at the Bottom of which is deposed a Sediment in Colour resembling pounded Brick or Bole Armoniac.

This Appearance is however feldom observed in any other but autumnal intermittent Fevers, and it is more so after several Paroxysms. In the Spring Intermittents, the Urine is commonly less red, and rather yellowish, a Cloud forms in the Middle, and deposes a white Sediment, which is a good Symptom.

Of the two Species of Intermitting Fevers we have just mentioned, the vernal ones are easier cured than the autumnal ones, which are attended with more troublesome Symptoms.

el Sint off recording to the Trick in

The Intermittents which prevail from February to July are called Vernal, and those that begin in the latter End of July, or Beginning of August, and cease at the End of January, or sometimes sooner, are called Autumnal Intermittents.

After the long and violent Heats of the Summer, if the Troops have been much fatigued, there will be many Autumnal Fevers, and of the most dangerous Kind, especially if the military Operations require camping in marshy Places.

In September and October, the Number of these Fevers is commonly very considerable, but there is great Hope to see that Number diminish at the Fall of the Leaf, especially if the Winds blow any Thing hard.

As there is a great Difference between the Spring and Autumn Fevers, and as the Method of treating them often differ much, we shall treat separately of each.

y favillable to such an ed or processor, and a

and have cashed and the last come

Make Sunt Witton

### SPRING INTERMITTENTS.

SPRING INTERMITTENTS are generally Tertians, very often of a kindly Sort; fometimes double Tertians, but seldomer than in Autumn.

We call that Fever a double Tertian, where a fresh Paroxysin comes on every Day, but the Fit is generally slighter on the alternate Days.

During the Paroxysm, it will be sufficient to take a large Quantity of any diluting Liquor, made agreeable to the Palate, but always warm, since cold Draughts would be hurtful.

The Patient may therefore drink of the Ptisan No 25. keeping himself quiet and in a moderate Degree of Heat.

The Fit most commonly goes off by an universal Sweat, which must be kept up by warm Drink, but ought not to be rendered excessive by too much Covering, or other Means that provoke Heat.

Just at this Time, viz. on the going off of the Fit or as soon as it is over, the Patient must have a Mess

a Mess of Broth with some Lemon Juice, or Cream of Tartar in it to make it acid.

The intermediate Days that are free from Fever, he may take Food somewhat more solid, viz. a little Meat, provided the Flesh be of young Animals: Beef will not hurt, so it be but tender, but all Sorts of Fat are to be avoided.

Nothing is to be eat near the Time that the Return of the Paroxysm is expected; the Nourishment then taken would load the Stomach during the Fit, and cause a bad Digestion.

Four Hours however before the Fit, the Patient may take fome light Broth. As in Spring Agues, the Paroxysms commonly anticipate the Time they should return at, Regard must be had thereto in the taking of Nourishment.

If the Day the Patient is without Fever be ferene, it will be proper for him to use a little Exercise: But not to Lassitude; he ought likewise to endeavour to seep rather more than usual.

It is to be remarked, that those Spring Agues often turn to inflammatory Disorders, especially in young and sanguine Subjects: Bleeding is therefore proper, especially if the Sick have a Redness in the Face, a violent Head Ach, or feel some Pain on the Side of the Breast.

If attended with frequent flatulent Eruptions, if the tongue is charged, a bitter Taste in the Mouth, or a light Vertigo, it will be proper to give him an Emetic.

Let him take four Hours before the Return of the Fit the Powder N° 26. or N° 27. if of a weaker Constitution, as soon as he shall have vomited, let him drink warm Water plentifully; he will soon throw it up, and vomit asresh; he then must repeat his Drinking, and go on in this Manner till what is to be thrown off his Stomach be diluted, and his Vomiting made easy.

After vomiting feveral Times, the Water drank commonly stays one Hour: After the Vomiting hath ceased, let the Patient have the Potion N° 24, and so wait for the Paroxysin, during which let him take the Ptisan N° 25. observing the Rules before laid down.

If he complains of Pains cross his Back, of Grumbling or Wind in his Bowels, if his Belly is swelled or hard, he must be purged in the following Manner.

Eight Hours before the return of the Fit, let him take the Purge N° 28. and fix Hours after he hath taken it, that is two Hours before the Fit, give him the Draught N° 24.

de this core an diffe

If the Symptoms that gave occasion to the Purge, or the Vomit, continue, these Remedies may be repeated, which however is but seldom the Case in Spring Fevers.

Observe, that sometimes Emetics do not evacuate only by Vomit, but by Stool also, and that Purges act likewise sometimes by Vomit. There is nothing to be seared when this happens, since the only Object of these Remedies is to evacuate the Stomach and Intestines. After the Bowels are thus cleansed, let the Patient take every two Hours a Spoonful of the Mixture No 29. drinking after it a Cup of Camomile Flower Tea. This Remedy must not be used in the Paroxysim, but only whilst the Patient is without Fever, nor should he be waked to take it.

This is the Method of treating Spring Fevers: and there is feldom a Necessity for using the Bark.

If after the third or fourth Paroxysm, ulcerated Pustules break out about the Nose or Lips, it is a good Sign, and the Fever quickly ceases: But this is not so sure in Autumnal Fevers,

It fometimes happens, though rarely, that after feven or eight Fits, the Spring Fever does not E 3 ceale, cease, nor even considerably diminish, and that on the contrary, the Fits become longer and stronger. This is particularly the Case, where the Patients are subject to sweat copiously, as soon as they are in Bed. Here the Bark becomes necessary.

Let him take every three Hours while the Fever is off, one of the Powders No 30. in some Wine.

This Method will foon compleat the Cure, and as in the Spring the Weather grows better every. Day, there is but little Fear of a Relapse.

The Property Could be the Country of the Country of

a se antique de los plessands de la company de la coloque la coloq

le tolk opher di bos Vergopap est panat il. Pank i kom komense bos kan i komense se se se se

And the said of the said of the said

growth business and the last

AUTUMNAL

## AUTUMNAL INTERMITTENTS.

THESE FEVERS are more obstinate than those in the Spring; and of these the worst Sort happen commonly after a very bot summer.

They are also harder to be known, for when they first begin, the Fits are so long, and the Returns so frequent, that they seem continued Fevers, with little or no Intermission.

Sometimes the Fever abates a little, and then returns in a few Hours, after a light Shivering. It is only when it begins to give Way its Character is first known; it then appears that the Diforder is a true Intermittent. These Fevers, which at first appear to be continued, often degenerate into Quartans.

Sometimes these Fevers, which in the Beginning did intermit, after long and redoubled Paroxysms, are changed into dangerous continued Fevers.

These Fevers are always bilious; the Stomach and Intestines are filled with putrid Matter, which must be discharged without Delay; to postpone it would be detrimental.

Let the Patient take the Emetic No 26. or 27. having Regard on this Occasion to what has been faid on the Subject of intermitting Spring Fevers.

If the Skin of the Face is tight and red, the Eyes inflamed, and the Heat great and general throughout the Body, a Bleeding must precede the Vomit.

On the contrary, if the Face is fhrunk and pale, and the Pulle not full, Bleeding would hurt, and must be refrained from.

The Vomit must be given in the Intermission of the Fever; or if it does not cease entirely, that Instant is to be chosen when it is least violent.

Sometimes also, in an autumnal Fever, it is necessary to sepeat the Vomit, that is, when the nauseous Sickness, the bitter Taste of the Mouth, and the Foulness of the Tongue, continue.

The Day the Patient takes no Vomit, let him drink the Decoction No 25. adding an Ounce of the Oxymel No 31. to every Pint of the Decoction.

After the first or second Vomit, the Powder No 32, is to be taken every four Hours.

This

This Method will commonly succeed in these Fevers; and if before, they were continued, they will become intermittent, so that there will be a confiderable Interval from one Paroxyfm to another, let the Patient then take the Mixture No 29. following the Directions given under the Head of Spring Intermittents.

The Food should likewise be the same as in these Fevers: Broths made pleasant with Juice of Lemon, or Cream of Tattar, roafted Apples or Pears, and well fermented Bread, ought to be the principal Nourishment. When he begins to recover a little more Strength, you may add fome flesh Meat, either Veal or Lamb: A little Wine taken moderately to recruit his Strength will do no Harm.

But as in Autumn the Days are continually growing colder, the recovering Patient must guard with Care against the Inclemency of the Weather, otherwise a Relapse is much to be feared.

color to locality with the Sand Sand State of Let him also for a Fortnight, on his Recovery, take the Quantity of a Nutmeg of the Electuary No 33. in the Morning falling, an Hour before Dinner, and an Hour after Supper.

and obtlicate Dispersa. It may be obtened, that When he has been a Month without Fever, give him the Pills No 34. in the Morning fasting, repeat them a second Time in eight Days, and a third Time in eight Days more.

If, after the Use of the Emetic, and other Remedies here described, the Fever should still continue, without any Diminution of the Paroxysms, and if the Patient grows weaker, the Use of the Bark becomes necessary: This happens more frequently in Autumnal, than in Spring Fevers.

The Powders No 30, must then be made Use of, as in Spring Fevers, and in a Fortnight be as gain repeated. S. Cathagains

If the Eyes grow yellow, the Sick feel great Anxiety at the Orifice of the Stomach, and the Urine is icterical, you must then (unless the extreme Weakness of the Patient forbid) leave off the Use of the Bark during the Space of a Fortnight; instead of which, for some Days give him two Spoonfuls of the Mixture No gig, every three Hours, till these Symptoms are abated? The Fever will return; but during this Interval, the Sick having recovered Strength, will better support it, and it foon will absolutely leave him.

If the Use of the Bark in this Case was obstinately purfued, it would bring on fome chronical and obstinate Disorder. It must be observed, that when the Fever has been driven away by the Bark, established the state of the state of the

the Pills No 34. are not to be made Use of, for commonly they bring it on again.

It sometimes happens, that these Fevers are from the Beginning attended with the worst of Symptoms, an unequal Pulse, a cadaverous Countemance, frequent Faintness, and cold Sweats. In some Subjects, these Symptoms are attended with a Cardialgia, or violent Pain of the Stomach, and in others with a sleepy Heaviness, sometimes so profound that the Patient can scarcely be waked.

In this Case the Bark must be immediately given; for it is to be feared another Fit could not be bore. The Prescription No 30, may be used.

If by this Means the Fever being suppressed, the Face becomes the Colour of Wax, and Anxieties, about the Orifice of the Stomach are complained of, let the Remedy No 35, be used in the Manner before explained.

the second of the real of reference and and are left I me or the Rod of a Forestell, by which be considered will be no Fear of a Rebusta.

easing the test of the common design of the state of the series of the common design of the c

The party state of the party of the second

QUARTAN

. 13550

commonly they bring it on agilt c

433

# QUARTAN FEVERS.

the Beginning granded this the week of himse HE first Thing to be observed is, that Bleeding is fearcely ever proper in thefe 

Let the Patient take the Emetic No 26. or 27. before the Fit comes on, following the Method prescribed in the Spring Intermittents.

In this Care the Bark stuff be immediately give

Before the next Paroxylin, give him the Purge No 28. and then the Quantity of a Nutmeg of the Electuary No 36. which he must leave off while the Fever Fit is on him.

If, after eight Fits, the Fever does not diminish, and the Patient grows weak, he must take the Powders of Bark No 30. observing what has been before faid on that Subject.

Were bearing the Col atto Way, and Ansieties.

Eight Days after the Fever has left him, the fame Powders No 30. are to be repeated, and a third and last Time at the End of a Fortnight, by which Means there will be no Fear of a Relapse.

In the intermediate Days of the Paroxysms, the Sick may be indulged with more Wine and a freer Nourishment, in this Sort of Fever, than in any other.

The

# THE JAUNDICE

A FTER a Fever, that hath continued for a Length of Time, and above all, after an Autumnal Fever, sometimes the Hypocondria remain hard and intense, either with or without a dull Pain; the Patient seels after his Meals an Anxiety, which sometimes is followed by Vomitings; the White of the Eyes grows yellowish; the Urine is tinged of an obscure yellow, and this Colour quickly spreads over the Superficies of the Body.

This Illness is also a common Consequence of bad nourishment; and the Soldier is most subject to its Attacks, when a Scarcity of Provisions hath obliged him to live on Meats of difficult Digestion.

Let the Patient take every three Hours four Spoonfuls of the Mixture N° 35. drinking after it four Ounces of the Decoction N° 37. which may any where be easily prepared.

Night and Morning let him have half a Drachm of Venice Soap in Pills, and his right hypocondrium be rubbed with Flannel every Morning, for a Quarter of an Hour, while fasting.

F

## JAUNDICE.

This Method followed some Days, generally renders he Base open, and procures Relief; but it much be construed till the Urine recovers its natural Colour, and the Yellowness disappears from the Eyes and It. Skin

If the Book is still costive after having used the Medicines six Days, you must give the Pills No. 14. In the Morning, and all that Day abstain from the Use of the other Medicines, which must be resumed the following Days.

Exercise is excellent in this Distemper, above all in open Air, if the Weather permits.

not domente a clis al alanill si

Farinaceous and glairy Food is to be avoided, and Broth wherein are boiled Chervil, Sortel, Lettice, Endive or fweet Success, is proper to be used.

Les the Preinc take every thirds Liours thur

Webs and Morning for him fare half a Dechar

214.1

Spacefuls of the Misseys Nº 353 delekissesiget it felic Ourses of the Decofflet 125 get, which have increased near the test of the parace.

Sur or wole allers and to the good sprandades

Geigeter of in Supr. - while fatting

## THE DROPSY.

WHEN the aqueous Part of the Blood is amaffed together, and retained in any Cavity of the Body greater or less, this Disease is called a Dropsy.

The Denominations of it are different, according to the Parts of the Body affected.

If the Water is detained in the adipose Membrane, and thereby causes a general Swelling, it is called Anasarca. The Swelling generally begins in the inferior Parts, and gains insensibly the whole Body; the Eyes are languid, the Face and Body sallow, the Urine in small Quantity, and the Patient never Sweats. On pressing the swelled Parts, it pits. The swelled Parts, and chiefly the Thighs and Feet are always very cold.

It is common enough in the Army for this Difease to succeed Intermitting Fevers of long Duration, especially in Autumn and Winter. The Soldier is also subject to it, when after having suddenly drank a Quantity of cold Water, he rests himself in a cold Place: It is also frequently the Consequence of a considerable Loss of Blood, either from Wounds, or Bloodings too often repeated.

F 2

When the Anafarca comes on after a long Intermittent Fever, Evacuations are not extremely necessary; but it is commonly cured by giving thrice a-Day, viz. in the Morning fasting, one Hour before Dinner, and one Hour before Supper, two Ounces of the prepared Wine No 38.

To finish the Cure, the Patient must be kept warm, either by the natural Heat of the Air, or by an artificial one: He ought to be well covered in the Night, keep to dry Food, such as Meat or Fish roasted; his drink should be little and pure, and as much Exercise be used as his Strength will admit of.

It will be found useful to rub the swelled Parts with warm Flannels, as often as you can. If the Patient voids Urine in larger Quantity, and begins to sweat in Bed, and the swelled Parts come down, it is a very favourable Sign.

After the Swelling hath disappeared, there remains such a Relaxation in the Parts as threaten a Relapse: This may be prevented by causing the Patient to wear his Cloaths streighter than usual, and rolling his Legs and Thighs with Bandages. Bodily Exercise in the open Air, and warm Weather does infinite Good.

In this Manner generally the Anafarca that fucceeds Intermittent Fevers is cured happily enough.

But

But when this Diftemper proceeds from other Causes, it is often more obstinate, and requires large Evacuations of the ferious Humour. Many Ways are tried to procure these Evacuations, but Experience shews the Remedy No 39, to be sure and efficacious. The Patient is to take a Spoonful of this in the Morning; or if, as it sometimes does, it brings on a vomiting, only half a Spoonful. The most common Effect is, however, only a simple Sickness.

The Urine after the Use of this Remedy comes away in large Quantity, and affords a confiderable Relief. It seldom purges; but if it should, does no Harm.

This Medicine is to be taken every Day, till all the Serofities are evacuated, and the Swelling of the Body entirely comes down. If the Dose is not efficacious enough in robust Constitutions, it should be infensibly augmented till it procures a large difcharge by Urine. When the Patient is on the Recovery, the same Regimen must be observed as has been just described. of the Renne

Sometimes the extravalated Lymph is accumulated in the lower Belly, and the Quantity increases fo much as to cause it to swell excessively. When fo, by pressing with one Hand on one Side of the Belly, and firiking on the other with the other Hand.

Hand, the Fluctuation is felt, and Water is found Can is, it is often more concerns, in bounded. Legistade la monagoria emil

When the Distemper is recent, it is cured often enough only by the Use of the Remedy No 39. but if the Flux of Urine does not come on in some Days, nor the Swelling of the Belly diminish, Haste. must be made to draw off the Water by the Puncture. The Operation is fure and fafe enough; but if attempted when the Disease is more inveterate, it is less efficacious.

It will be proper, as much as possible, to draw off all the Water altogether at once: Which may be done fafely, by streightening the Patient's Belly with a Bandage, by little and little, and more and more, as the Water runs off: The Faintings and other Accidents will, by this Means be avoided.

The Using affice the I've of this Roughly concer

the Shillestide's comes down. If the Link lance After the Evacuation procured by the Puncture, the Belly must be bound tight with Rollers, and the Patient observe the Diet we have described above. governo do se tigan be regard sie al ses

The Use of the Remedy No 38. will be very proper here. Sometimes the Belly swells again, and the Puncture is then to be repeated.

As it sometimes, though but rarely, happens, that the Swelling of the Belly is caused only by Wind,

Wind, and not at all by a Collection of Lymph, the most scrupulous Attention must be used in observing what the Case is, because in this last the Puncture is scarce ever of Use, and, on the contrary, generally accelerates Death.

This last Disease is called Tympanites, or Dropfy of the Belly.

- ed as in the Dropfy.
- The Belly appears elevated before, and depressed on the lateral Parts.
- 3. When the Belly is struck, the Motion of Water is not felt, but it gives a Sound somewhat reifembling that of a Drum.
- 4. Though the Sick lies on either of his Sides, the Form of the Belly remains the same; the Skin of which is white, extended, and elastic.
- often precede this Distemper.

connect tie down, but is chipped as the op in his

This Diffaile is known by the presenting Canfer.

6. The Body of the Patient is lighter than in a Dropfy of the Belly, where the Collection of Water confiderably augments the Weight.

nigu

Upon the whole, this Diftemper is more dangetous than the other, and is often mortal.

O - Will Make the bate for W

The Cure may be attempted in the following Manner. Let the Patient's Belly be rubbed with Flannel, for a Quarter of an Hour at a Time, twice a-Day; and after each Friction, let it be anointed with the Liniment No 40. and for feveral Days let him at Night take the Powder No 41.

If the Winds begin to pass by the Anus, and the Belly grows less, there is Room to hope a Cure.

It happens fometimes, that the Lymph collects in the Cavity of the Breast; and it has been often observed, that Soldiers are subject to this Kind of Dropsy, when, after having been heated by Work, and all in a Sweat, they suddenly expose themselves to the Cold, and greedily drink cold Water.

This Disease is known by the preceding Causes, by the Dissiculty of Breathing, especially when the Patient begins to sleep, and by a dry Cough; he cannot lie down, but is obliged to sit up in his Bed, his Body bent forwards, and has commonly his Feet swelled at the Beginning of this Distemper.

It is observed, that the Breast is sometimes disengaged, when the Swelling of the Legs and Thighs become very considerable; and that, on the contrary, the Breast becomes more oppressed, when this Swelling suddenly disappears.

This Sort of Dropfy, especially if not inveterate, is often successfully cured by the Use of the Remedy No. 39. Decide of About 1 and 1

If this does not answer, nothing remains, but the Puncture; but we find by Experience, that this is but a doubtful Means, and not always attended with Success.

the Pargue to baded by a piteitous glairy Matter, of light Condia, field as No. 37, thould be given, and the hard the fauth Regulars while red as preferibed in fractulators Freeze.

so When the Latin order door verifier his him, take the Man take the Court of the Man take the Mount of the Man to the Manny of the Day he take the Many him to the Many him him to the Many h

COLERAMOREU

VOMITING.

TT & hidden and immoderate Recumina of A the Humours, bods upwards and downstant

Locgi

### VOMITING.

become your confidentials on a property

OUR present Inquiry does not regard those Vomitings that attend other distempers, as Fevers, nephritic Cholics, &c. but only those occasioned by bad Food, and a loaded Stomach.

The furest Remedy, in this Case, is to cause a large Quantity of warm Water to be drank, to render the Vomiting and Evacuation of soul Matter easier.

If after this there remains any Nausea, or that the Tongue be loaded by a pituitous glairy Matter, a light Emetic, such as No 27. should be given, and the same Regimen observed as prescribed in Intermittent Fevers.

When the Patient has done vomiting, let him, take two Spoonfuls every three Hours of the Mixture No 42, and on the Evening of the Day he takes his Vomit, let him have the Draught No 24.

#### COLERA MORBUS.

IT is a fudden and immoderate Evacuation of the Humours, both upwards and downwards.

Though

Though this may happen, at all Times of the Year, in consequence of overcharging the Stomach, and a Crapula; yet it is most frequent towards the End of the Summer, and Beginning of the Autumn. It is often occasioned in Summer by eating too much Fruit, drinking putrid Water, or a great Quantity of new and sweet Wine, called Must.

The Violence of this Disorder is such as to reduce the strongest Man in a few Hours, and sometimes to carry him quite off in the Space of twenty-four Hours.

The Thirst is commonly violent, the Anxiety great, the Pulle quick, small, and often unequal. The Patient has cold Swents; his Pace is wan and cadaverous, and his Extremities cold.

The Thighs or Hands are attacked with Spains, and sometimes both together; all these Symptoms are soon followed by Convulsions, and by Death, unless prevented by the most speedy and efficacious Remedies.

Every Emetic and Purge is to be avoided, for the most gentle are hurtful in this Case. The Patient must be continually taking Chicken or Veal Broth, made so light as scarcely to have the Taste of Means or instead of these, Panada. A Glyster of either of these Drinks is likewise to be given, in order order to discharge all acrid and irritating Matter from the Intestines.

After this Method has been continued for three or four Hours, a Spoonful of the Mixture No 43- is to be given every half Quarter of an Hour, till the Vomiting and Looseness are either stopped, or considerably diminished.

When the Patient begins to be somewhat better, it will be sufficient for him to take only everythree Hours a Spoonful of the same Medicine, till he has taken it all.

Though the Vomiting and Looseness should by this Means be entirely stopped, let the Patient still continue, for four Days, to take three Spoonfuls of the same Remedy N° 43. every Night and Morning.

The best Food, on these Occasions, is Veal Broth with Rice in it, taken a little at a Time, and often.

When it happens that the Sick lies long without Help, and has had these Evacuations for several Hours, and consequently is grown very weak, and especially if he seeds Spasms in his Thighs or Hands, you must immediately recur to the Mixture No 43. in the Manner above described.

Broth, made to hight as fear sky to have the Taile of Mylett A inflead of thefe, Paneda. A Glytlet of thete Drinks is likewife to be given, in order

#### ADJARRHÆA

de this a sa state

WHEN the stools are liquid, and more frequent than usual, the disorder is called a Diarrhoa.

The pains and gripings of the belly are not violent in this diftemper; which by this principally is diftinguished from the Dysentery, which we shall treat of afterwards.

As the Diarrhea sometimes serves for the evacuation of bad humours, it follows, that it is not always hurtful, and that it may even be advantageous.

It is of service when it does not affect the strength, but, on the contrary, renders the Body more light and alert. On the other hand, it is hurtful when it brings on Languor and Weakness.

The Diarrhæa, that at first appeared serviceable, may become detrimental, by its too long Duration, viz. if it lasts four or five Days; for then the Body is worn out by too long a Flux, the Intestines excoriate, a lively Pain of the lower Belly, with gripings, succeeds, and the Diarrhea degenerates into a Dysentery.

W

When the Diarrhea requires any Remedy, you may give the Powder N° 44, and at Night the Potion N° 24. Veal Broth with Rice, and Millet boiled in Milk to a Thickness, are proper Food for the Sick.

If the Diarrhæa does not stop in two Days, the Powder No 44. and Potion No 24. must be repeated, and again in two Days, if it should so long continue.

of double the base

isman (bed) look longuage

To avoid a relapse, let the patient, on his recovery, for four Nights successively, take the Bolus No 45.

Care must be taken that he be well covered and cloathed, and guarded against the Injuries of cold Air.

Server Serve alies of the server

liefere web grouping for Secretaries for Alberta

The

#### The DYSENTERY.

ADIARRHÆA too long neglected often degenerates into a Dyfentery.

But it most commonly happens, that it is not preceded by that Disorder; and in the Army it reigns during the Heat of Summer, and the Beginning of Autumn.

A Flux of the Belly, attended with violent gripings, and very painful strainings for Stool, is called a Dysentery.

The Stools are not always accompanied with Blood, as feveral Physicians have pretended, who therefore have called this Diforder the Bloody Flux.

Nevertheless, the focal Matters are often reddish and bloody, especially if the Disorder has lasted any Time.

This Distemper prevails chiefly among the Troops. The Causes that produce it are as follow:

The Bile grown acrid by the great Heats and the Fatigues of War; especially if the Soldier, G 2 when when heated, suddenly exposes himself to cold Air, or sleeps in his Cloaths, soaked with Rain: for which Reason it is often brief in Places, where cold Nights succeed to hot Days.

Stagnating or Marsh Water for common Drink.

Meat, or Fish, beginning to be tainted, musty Bread, or Bread made of musty Corn for Food.

Sure and reiterated Observations convince us, that Summer Fruit scarce ever cause the Dysentery; but the Excess of them may do Hurt.

This Distemper proceeding from the Causes just described, soon infects a whole Army: The healthy soldiers are more especially infected by the putrid exhalations of the second Matters, if they use the same Bog-houses as the Sick.

This Point must therefore be carefully attended to, when the Dysentery prevails among the Troops: It would be proper to dig deep Trenches to serve for Necessaries for the fick Soldiers, to cover several Times in a Day the secal Matters with Earth, and to have other Trenches reserved altogether for those in Health.

To change the Camp often would also be a proper Means to stop the Progress of this Disorder; and by what has been said relative to the Causes, it will appear necessary to take all possible Care to avoid them.

The Manner of treating the Dysentery is as follows:

If the Sick is of a fanguine Habit, and has great Heat all over his Body, or much Fever, he much lose eight or ten Ounces of Blood from the Arm; but these Symptoms are rarely met with.

As the Dyfentery is feldom attended with a Fever, and then Bleeding is of no Use, it will be sufficient to give the Patient the Powder N 46. in Wine.

After the first Vomiting occasioned by the Powder, he must drink warm Water with a little Honey in it: This will excite fresh Vomiting, after which he is still to be plied with warm Water all he brings it up clear as he drank it.

After the Patient shall have rested two Hours from his last Vomiting, you may give him some small Slices of toasted Bread, soaked in sour Ounces of cold Wine; and to make it pleasant, a little Cinnamon and Sugar may be added. At Night let him take the Pill No 47.

The same Remedies are to be repeated next Day; and if the Disorder is not entirely, or much abated, the third Day again. But if the Diffemper is confiderably diminified, it will be proper to leave a Day's Interval between the Use of these Remedies, before they are given a third Time.

Experience has flown, that instead of the Powder No 46, that No 48, with the Pill No 47, at Night, has been given with good Success. This must also be repeated three Times, leaving the Interval of one Day, except the Dysentery should stop sooner. If the Remedy No 48, he too weak for robust Constitutions, the Dose may be increased to ten or twelve Grains. After these Evacuations, the Patient may for several Days take at Morning, Noon, and Night, one Drachm of the Electuary No 49.

His Drink must be two Parts in three of Barley or Millet Water, and one of new Milk; and this is to be used in large Quantities.

For Food, Milk thickened with Barley, Oats, Millet, or Rice, may be given; and when the execsive Stench of the Stools is somewhat abated, Meat Broth thickened with the same Pulses.

But if the Malignity or Duration of the Distemper, should have, in a Manner annihilated the Strength of the Patient, evacuating Medicines must be laid aside, as he is already but too weak.

This

This is known to be Case, by the Violence of the Griping and Strainings, by the varillating small Pulse, by the pale Countenance, by the nauseating all Food, and by an unextinguishable Thirst.

Let the Patient take every Hour one Ounce of the medicated Wine No 50, and Night and Morning the Pill No 47.

As the bad Symptoms begin to disappear, and Strength to come again, let the Powder No 44. be taken in the Morning, and in the Evening the Pill No 47, to be repeated thrice, with one Day's Interval, if the Disorder be not over sooner.

For some Days after, let the Patient take one Drachm of the Electuary No 49. thrice a-Day.

Sometimes the Intestines having been exceriated, by the frequent Passage of acrid Matters, the Patient is very much incommoded with a continual Inclination to go to Stool, the he voids little or nothing: In this Situation he is to have the Glyster N° 51. and to retain it as long as possible.

If, after the Evacuations, there should remain such like Pains in the lower Belly, the Patient would receive great Relief, by swallowing every Day, an Egg boiled soft, with a little fresh Butter.

#### INFLAMMATION of the INTESTINES.

A N Inflammation of the Intestines, a very dangerous Distemper, often arises from the same Causes as a Dysentery.

It is discovered by a violent Pain of the lower Belly, which often increases upon the Touch; by the Swelling of the Belly, by Vomitings, and by Constipation. These Symptoms are attended, at the same Time, with an acute and continual Fever, and a violent Heat: The Pulse is hard, the Urine clear, and of a bright red, and there is a studden Prostration of Strength.

If the Symptoms are violent, most commonly Death is soon-the Consequence. Before the Patient expires, the Pain ceases; but the Extremities become cold and livid, the Face cadaverous, and Pulse small, quick and unequal. All these are Signs of approaching Death; although the Sick, and those about him, are apt to draw an happy Omen from the Cessation of Pain.

Copious Bleeding is immediately to be had Recourse to, and repeated boldly if the Pains do not go off, or remit considerably; or if they return again. Let the Patient take the Glytter No. 52, three or four Times a-Day.

His Belly must be constantly fomented with. Flannels steeped in the Fomentation No 12.

The Caul of an Animal fresh killed sometimes produces a good Effect.

Let the Patient take every Half Hour a warm Cup of No 53.

It is a good Sign if the Pulse becomes equal, and remains so, if the Pain diminishes, the Sick break Wind downward, and the Glyster bring away with it see Matter.

Sometimes, notwithflanding feveral Glysters, the Belly remains obstinately constipated: In this Case the Fumes of Tobacco injected through the Anus have produced very good Effects.

The Drink is to be warm Barley Water, and light Broths the whole Nourishment, until three Days after the Disorder shall have entirely ceased.

And even then it will be necessary he observes an exact Diet for some Time, less the Intestines, irritated anew by acrid Food, occasion a Relapse.

the Rana out of the Ableckeis not leng-

This Distemper is indeed so violent, that if it does not quickly give Way to proper Application, it degenerates immediately into a mortal Gangrene. But by the careful Use of the above recited Remedies, one may hope to resolve the Instammation of the Intestines.

If this Method has begun to be used too late, if the Distemper lasts, without growing worse, three or four Days, and a dead Pain succeeds to the acute Pain of the lower Belly; if at the same Time the Patient seels an unusual Heaviness, and has wandering Shiverings all over his Body, it is certain there is an Abscess forming.

In which Case he must have the Fomentation No 12. constantly applied on his Belly during the Day, and at Night a Plaister of Labdanum.

If the Abscess seems to be ready to pierce externally, which may be if the Intestines adhere to the Peritoneum, it must be opened to discharge the Pus. But this Case seldom happens.

If the Abscess breaks in the Cavity of the lower Belly, the Consequence is much to be feared, unless the Matter can be drawn off directly, which is very difficult to be done: Nor is it easy to judge of the existence of this Case, because the Quantity of Matter from out of this Abscess is not considerable

INFLAMMATION of the INTESTINES. 71 derable enough to cause any remarkable Swelling of the Belly.

The Pus is more frequently evacuated by the Anus: The Glyster No 52 repeated several Times after Suppuration is perfected, facilitates its Course. By the smoothing of the internal Coat of the Intestines the Evacuation of the Matter becomes eafier that Way.

After the Pus is evacuated, whether alone or with the Excrements, the Patient must drink plenfully of the Decoction No 16. sweetened with Honey, and is to take the Powder No 18. three Times a-Day.

Let Endive, Lettice, Chervil, or such like tender Herbs, be boiled in Broth for his Food; but it must be strained thro' a Sieve, lest any thick substances should collect into a Mass in the Intestines.

Let him continue this Method for three Days after the Pus shall have ceased passing through the Anus; and by Degrees he may return to his former Way of Living.

THE COUNTRY OF PAIR COUNTRY IN

the state of the s

## PHRENZY.

A PHRENZY is a continued Delirium, attended with an acute Fever. It is thereby distinguished from the Delirium, which is sometimes observed in the Height of intermitting Fevers, and which sinishes with the Fit.

An extreme Heat, and a violent inflammatory Head-Ach, commonly precede the Phrenzy: The Eyes and Face are red; the Patients, when asked Questions, answer with Ferocity; they pluck the Knap of their Cover-lids.

The more frequent Causes of this Distemper are the violent Heat of the Sun, to which the Soldier is exposed, especially if bare-headed, sleeping in that Condition, long Watches, extreme Emotions of Anger, Excess of Wine, Brandy, or other spirituous Liquors.

In this Distemper the Pulse commonly is quick, and Respiration strong, and not frequent.

A Phrenzy is very dangerous, and often occasions sudden Death, for it is a true Instammation of the Meninges, and sometimes even of the Brain itself.

Greenish

Manager Serie

Greenish Vomitings, frequent Spitting, Shiverings, crude, aqueous and pale Urine, Convulsions, and no Thirst, are bad Signs: Bleeding Piles, a Flux of the Belly, a copious Hæmorrhage of the Nose, give Relief to the Patient.

Pain of the Breast, or of the lower Parts, is good in this Illness: A strong Cough coming on, sometimes also gives Ease.

foreset.

Bleeding here is effentially necessary: It should be large, and chiefly in the foot: If the Fever and heat continue, it must be repeated: And it will be right, after the first bleeding of the Foot, to open one of the Jugulars.

The Bleedings must be repeated; till the extreme Heat and the Violence of the Delirium are abated. While the Patient does not sleep, he must take every Hour a Cup of the Remedy No 54. warm.

For his common Drink, the Decoction No 25. may be used, and Night and Morning the Glyster No 11. be given.

If the Hæmorrhoids swell, let Leeches be applied.

It will be proper to have the Patient's Head, and to make him rinse his Mouth often with warm H Water: Water: Let a Compress dipped in Oxycrate, or Vinegar, and Water, in equal Quantities, be applied on his Forehead: Fresh and temperate Air is most proper for him, and he ought to be kept sitting upright in his Bed, with his Head raised as much as possible.

It would be also proper to make him rise twice a-Day, to set him in an Arm-chair, and bathe his Feet in warm Water. After the Night-bathing, the Paste No 55. should be applied to the Soles of his Feet, and lie on till Morning.

During the whole Time of the Illness, he must be confined to plain Barley - Water, or Water-Gruel.

WHALLS I

begins sensibly to grow less, and the Delirium to wear off, but the Patient can get no Sleep, let him have the Emulsion No 17. at Night, with the Addition of an Ounce and a Half of Sirop of white Poppies.

But Anodynes must be carefully avoided, while the Illness continues in its force.

At the Beginning of this dangerous Distemper, all the Remedies here recited must be vigorously employed: But when the Heat and Delirium are considerably diminished, Bleeding and Glysters are no longer necessary, the Drink No 25. will be sufficient, and the Nourishment may be a little stronger.

Notwithstanding the Diminution of the Symptoms, it happens often enough, that the Delirium does not entirely give Way: But commonly it decreases insensibly, especially is several Times a-Day, and as much as his Strength will permit, the Patient is made to sit up in an Arm-chair with his Body upright.

Ber Berglinsberin Restland in 1821king in plotser, neisten ing Seland entrologie wasere entropy their college Search is to redichers co after an inel verenige in the vern Dank neg

The distribution of the control of t

Type of the delegant was the selection of the selection o

T

H2 HÆMOR-

the sale by but the

The array of the property and a party larger

water the point and the location of the principle on

#### HÆMORRHAGE of the NOSE

A S Bleeding of the Nose is a pretty common Symptom in ardent Fevers, almost always giving Relief, and sometimes even curing the Diftemper, it is easily apprehended, that it must not be too lightly stopped.

But sometimes the Bleeding of the Nose is so violent, both in sick Persons, and in those who are in Health, the Bodily Strength is so wasted, as to bring on total Faintings, so that even Death may ensue.

In this Case, the too violent Hamorrhage must be stopped. To judge when this ought to be done, requires a little Attention to the following Considerations.

While the Pulse keeps full, and the Heat of the Body is every where equal, even to the Extremities, and the Lips and the Face keep their red Colour, there is no Fear from the Hæmorrhage, was it even violent.

But when the Pulse begins to undulate, and the Face and Lips grow pale, it is Time to stop the Bleeding.

within man, and also List less in the Nove, where

the Oak is also an efficacions The Means of stopping it are, applying Bandages to the Arms and Thighs of the Patient, because the Veins being thereby compressed, the reflux of the Blood to the Heart is in less Quantity. The Hæmorrhage stopped, the Bandages are not to be loofened all at once, but fuccessively one after another, in such a Manner as to leave the Space of a Quarter of an Hour between the loofening of each Bandage.

If, by the Application of the Bandages in the Manner described, the Hæmorrhage is not stopped, or if it comes on again on taking them off, the following Method is to be used.

Let a Tent of Lint imbibed in the styptic No 56. be put up that Nostril from which the Blood came. If some Lint dipped in the Styptic be wrapped round a Quill, it will be eafily introduced up the Nose: At first, for about Half an Inch, it must be put up horizontally, then raising the Quill insensibly, it is to be pushed gently, and by this Means the Lint introduced as far as possible without hurting the adjacent Parts, Afterwards, by gently compressing the Nostrils, the Quill may be H 3 with-

#### 78 HEMORRHAGE of the NOSE

withdrawn, and the Lint left in the Nose, where it is to remain for a Day or two, till it falls out of itself.

The agaric of the Oak is also an efficacious Remedy for stopping the Bleeding. Some of the Powder No 57, may be blown thro a Quill into the Patient's Nostrils.

see March ringe from a the following the seed of the green of the green of the following the seed of the control of the seed o

Minister Caleriosa, in I mentioning in no single see, or it is comes on again on taking their off.

enter a species enter estate a maiore estate a monte estate enter enter

y comprehend the North the top one to

the following Methodis' to be tifed.

the postulation of a single

which proportions with the main of god

A CON-

1 . Pontil and Tear

# A CONTINUED FEVER

A FEVER, that lasts from the Moment of the first Access, without Interruption, to the End of the Disease, is called a Continued Fever.

The principal Causes of Fevers of this Sort in an Army are, excessive Fatigue, and extreme Lassitude, which is its Consequence, especially during the Heat of Summer, if the Soldier is under the Necessity of enduring Thirst, or drinks too much Spirituous Liquors.

For the most stuid and lightest Parts of the Blood being by these Means lost; what remains, grown more thick and acrid, is in a State to occastion great Disorders, and above all inflammatory ones, because the Mass of Humours so thickened is now greatly disposed to Inflammation.

When a Fever of this Sort produces a topical Inflammation, the Disease takes its Name from the Part affected: For the Pleurisy, Peripneumony, Phrenzy, Sore Throat, (Quinzy) Inflammation of the Intestines, are often preceded, and always accompanied with a Continued Fever.

210 1

But when it happens that some of the above recited Causes occasion this Fever, and that it affects no Part in particular, it is called simply a Continued Fever.

This Fever is known by the Causes that precede it, by the Vigour of the Time of Life, and a hot and sanguine Constitution, by a hard and quick Pulse, and chiesly by its extreme Heat, which burns, as it were, the Fingers of him who touches it. The Urine is red, thick, and turbid, the Tongue dry, the Thirst great; often an intolerable Pain of the Head, and an obstructed Respiration.

This Disease, always dangerous, is more or less so according to the Violence of the described Symptoms.

Large Bleedings are necessary in the Beginning, which are to be repeated till the great Heat and Dryness of the Tongue begins to abate. Barley-Water is the common and proper Drink; but to every Pint should be added an Ounce of the Remedy N° 31. of which let him take largely; let there be given him every two Hours a Cup of the Decoction N° 54. and twice a-Day the Glyster N° 11.

This Method is to be continued till the Decline of the Diftemper, which is known by the Diminution of Heat, of the Quickness of the Pulse, and of Thirst; by the Humidity of the Month and Tongue, by the Urine not being so high coloured, and by the Sediment it then deposes. The same Regimen to be observed as in the Pleuristy.

As the Disease grows milder, the Decoction No 25, will be sufficient for common Drink, and let the Diet be gradually augmented till he is quite well.

It is highly necessary to observe, that there is also another Sort of Continued Fever without an inflammatory Thickness of the Blood, but rather occasioned by a putrid Dissolution of the Humours. This last Sort is much worse, and more dangerous than the other, and very often this Fever proves contagious.

This happens most frequently, when, during the great Heats, the Army is encamped in marshy Places; for then they respire an Air corrupted with bad Effluvia. This Sort of Fever prevails also very much, where many Men, even were they healthy, are lodged together in a narrow Space, where the Air cannot be renewed often enough. Ships of War and Hospitals, where the Sick and Wounded are much strengthened, are

frequently visited by these Fevers, especially if the Air cannot be refreshed often enough; because the Air then to be respired is so corrupted by the Essure from the Bodies, the Stink of the Excrements, and the Putridness of the gangrened Parts, that it engenders a very bad, and truly putrid Fever, which soon grows contagious. It is therefore sometimes called the Jail or Hospital Fever. Its particular Symptoms should be exactly described, thereby to know this Distemper.

It begins by a Shivering, followed by a Heat, but not violent; foon after, the Shivering again, after which the Heat, and so the Shiverings and Heat alternately.

Total Loss of Appetite, Sleep disturbed, and without Refreshment, a heavy Pain of the Head, effecting especially the anterior Part: The Pulse is almost as in a natural State: The Skin is not always very dry: The Sick languish on in this Manner some Days, without being able to attend their Business, yet without being obliged to keep their Bed. The Tongve is seldom dry; it is more commonly soft, most, and covered with a Sort of Crust of yellowish Green. The Patient dozes much, sleeps little, and seems quite absorbed in prosound Reveries: In the Progress of the Illness, comes on a Trembling of the Hands, Hardness of Hearing, and Dimness of Sight; the Pulse begins to grow feeble, and the Parient

is defirous of Cordials and Wine. Towards Night all the Symptoms grow worse: Lastly, at different Times during the Course of this Diftemper, there appear purple Spots of an irregular Figure. is comment refunctional Merchan con-

The following Symptoms are confidered as mortal: A fudden Proftration of Strength, Weakness of the Sight: The Posture of the Sick, stretched on his Back, and drawing up his Knees to him; reiterated Efforts to get out of Bed, black Aptha, livid Petechia, and Stripes also livid, refembling the Blow of a Whip spread over the Body; the Flux of the Belly, with Lead-coloured or blackish Stools, weaken the Patient more and more! " I a first said as land a fa to first.

culton Whose that be mad vernice that which as to Deafness is not a bad Symptom in this Distemper. Nay it has been observed, that Patients on their Recovery are apt to grow deaf, and that fometimes they have an Abicess in the Conduit of the Ear.

Bilious Stools, thick Urine, a moist Tongue, are good Omens, especially if the Patient keep up his Strength. () se immi

A Number of little red Puftules, or white and elevated miliary ones, are good, if at the fame Time Expectoration is easy, and the Urine deposes a thick Sediment. Lastly, it is counted a good Sign, when an easy Sweat comes on, and relieves the Patient, or the Parotids swell, or there appear white Apthæ.

As the Causes here recited indicate that all Things tend to Putrefaction, and that the Strength is extremely exhausted, Bleeding can seldom be used, unless in replete Habits, and that but once: for large Bleedings abate the Strength immediately, and occasion a Delirium. It is very necessary that the Air be often renewed.

a much of should between a mid of

held thought on separation with the finish

In case of a Nausea, or Weight felt about the Region of the Stomach, or that the Tongue be covered with a yellowish green Crust, the Patient must take the Emetic No 27, and after the first Effect of this Powder, let him drink plentifully of warm Water, that he may vomit easy, which is to be repeated, as has been said in the Article of Intermittents.

The Evening after the Emetic, let the Patient take the Bolus N° 58. and drink after it fix Ounces of the Whey N° 59. If Milk cannot be procured, the Decoction, N° 25. may be substituted in its Room, observing to add two Ounces of Wine, and Half an Ounce of Oxymel, to every Pint. This Whey or Decoction may serve for common Drink, especially as the Sick are fond of vinous and comforting Drinks, and that these Drinks are suitable to this Distemper.

Let him take every fix Hours the Powder No 60. with fix Ounces of the above Whey or Decoction.

Death, preceded by great Anxiety and Convulfions, is often the Consequence of extreme Languors, when the Purples disappear, or the miliary Spots strike in: In which Case a Spoonful of the Mixture No 61. is to be immediately given, with three Ounces of the Whey or Decoction No 25: to be repeated every three Hours till he finds Ease, and the Purples appear again, or the Miliary Spots rife; after which the same Remedies are to be continued every four Hours only. If, in Consequence of this Method, an equal Breathing Sweat comes on through the whole Body, the Patient finds great Ease thereby. If the Belly is constipated, let the Glyster No 52. be given.

It is proper to discharge the Patient from out of the Hospital, as soon as they begin to recover, that they may breathe a purer Air; otherwise a Relapse is much to be feared, which is seldom or never got over.

penify and extude an borrel strack + the Teah

damen violent Humaning with vary used concid-

and he Limbs watch increase in the high traffic and

grow sellow, then birth and cuthan.

The appropriate party of the propriate that the

The citemate the transmit of course of many citement

Death, placeded by many while and Cometfrom it court the Confinence of extreme field-

## The SCURVY.

HIS is a common Distemper, and of dif-A hoult Cure, more especially in Sieges and unhealthy Places, where fometimes the Troops are obliged to take up their Winter-quarters.

the Louis of he finds It begins by a Numbness of the Limbs, with an unufual Lassitude of the whole Body: After walking, the Limbs and Muscles feel quite fatigued, and, as it were, broken. In the Increase of the Distemper, Respiration becomes short and difficult; fometimes the Thighs swell; at first the Face is pale, then grows brown, and the Skin is stained with Spots of different Colours, the Mouth begins to fmell, the Teeth loofen in the Sockets, the Gums swell, itch, grow painful, and bleed on the least Touch; lastly, wandering Pains affect different Parts of the Body.

In the Progress of the Distemper, the Gums putrify and exhale an horrid Stench; the Teeth grow yellow, then black and carious. Sometimes happen violent Hæmorrhages; very bad conditioned Ulcers break out, especially on the Thighs; the Patient feels violent and painful Shootings in all his Limbs, which increase in the Night, and the Body is covered with black Spots. At this Period the Distemper suddenly grows worse, Fevers come on of different Kinds, every Thing grows quickly putrid, and mortal Hamourhages break out, of the Mouth, the Nose, or about the Anus; the Viscera putrify; and Faintings succeed, which are soon followed by Death.

This Distemper is frequent in Winter-quarters; from the following Causes.

Series were associate actional Series

Noisome Vapours, arising from marshy Grounds and stagnating Waters, Inaction, Scarcity of Greens and Vegetables, drinking of corrupted and stagnating Waters, the Use of salted and smoaked Flesh and Fish, and of Cheese too old and acrid; damp and low Lodgings, and not being open to the Course of the Winds.

Fear and Sorrow also occasion this Disorder, and increase it in those already attacked with it. By that, and by bad Food, it often makes such Ravages in besieged Places.

Experience shews, that the Humours in this Disease are not only putrid and acrimonious, but also condensed.

Therefore, in the Cure, Care must be taken to attenuate the Viscosity of the Humours, and to prevent or correct Putrefaction.

We are to lend all the Affistance of Art to prevent, or avoid the Causes of this Evil, and thereby preserve the Soldier from the Ravages of this Distemper.

First, by correcting the impure Waters. This is done by mixing two Ounces of Vinegar, and two Ounces of Brandy, to every Pot of Water. For Want of these, some Slices of Calamus Aromaticus may be steeped in the Water. This is a Sort of Reed, very common, growing almost every where, in such low, marshy and damp Grounds, as are most subject to the Scurvy.

Strong Purges, Vomits, and Bleeding, do no Service in this Diftemper.

But as bad Nourishment is one Cause that produces the Scurvy, the Stomach and Intestines must be cleared, and evacuated, which is easily done, by gentle and reiterated Purges, such as No 34, to be taken three Times, with the Interval of a Day between each.

The Food should be Broth, with Chervil, Sorrel, Spinage, Lettice, endive, Succory, Cabbage, especially Red Cabbage, young Nettle Buds and Tops, or any other Sort of tender Herbage, boiled in it; the Preserence to be given to those easiest to come at.

Fruit

Fruit quite ripe, used moderately, always produces a good Effect: But if neither Fruit nor Greens can be procured, the Patient must have his Broth with Barley, Oats, or Rice; he may eat likewise a little Veal, or Fowl, but it must be moderately.

After the Use of light Purgatives, Antiscorbutics will be proper, but which are to be varied according to the different Constitution of the Patient.

If he feels himself cold, his Face pale, his Legs swell, and his Thirst is not great, let him take two Ounces, that is about a Tea-cup of the Decocion No 62. thrice a-Day.

If he is hot, his Pulse feverish, his Thirst great, his Breath bad, his Gums bleeding and half putrid, the Decoction No 62. is not so proper as No 63. to the Quantity of three Ounces thrice a-Day.

Ripe Fruits, and roafted Apples and Pears, eafy to be procured, are also very proper.

77 H and I shall in about

The Remedies are to be continued a great while. When the Limbs move easier, and the Pain diminishes, the Distemper grows better, and then Exercise and good Food will be sufficient to I 3 compleat

compleat the Cure. To carry off all Relicks, it will be proper, on Recovery, to take fifty Drops of the Elixir No 64. in Wine and Water, equal Quantities, thrice a-Day.

and a second of the second of

Altho' it be certain, that when the Distemper is at an End, the Symptoms ought likewife to cease; yet it is no less true, that after the Scurvy, we often fee those who have been attacked by it subject to Ulcers of the Gums, Lips, inside of the Cheeks, and of the Palate, which foon spread, and corrode these Parts, and in a little Time turn These Ulcers often deceive those to a Gangrene. who do not rightly understand them: They appear in Form of white or yellowish Spots, red and inflamed round the Border, and often very painful. A great Stench accompanies them, and the Spittle, which comes in Plenty, is also of a bad Smell. This Evil requires an immediate Remedy, otherwife all would foon be infected with a gangrenous putrid Humour, the Teeth would fall out of the Sockets, the Jaws would be affected, and entirely corrupted.

But this Diforder is eafily got the better of, by touching the Parts lightly and often in the Day with a little Lint dipped in the Preparation No 65. Little Compresses, imbued with the same, may also be applied between the Gums and Lip, and renewed from Time to Time.

n correspicac

Care

Care must be taken not to rub the Parts affected too much, as is the bad Custom of some, for the Evil and the Pain is thereby augmented.

If the Stench is great, and the Ulcers extend themselves rapidly, the Quantity of Spirit of Sea Salt is to be augmented, till you get the better of the gangrenous Corruption.

AND ENGINEER OF LONG POINTS AND COME TO SERVE TO

e a contrata de la companya de la co

spinis it is a little with the control with the

## The GANGRENE.

to se relative of to the Parts allegated

As Mencion has been made of the Gangrene, it may be here proper to take Notice, that the Bark taken internally is a most efficacious Remedy for this Distemper, whatsoever Part is attacked.

The Patient is to take every four Hours one of the Powders No 30. till the Gangrene begins to feparate from the found Parts, and a good Suppuration comes on: At which Time it will be fufficient for him to take them twice a-Day till the Ulcer is mundified.

The Bark is equally proper, when the Scorbutic Ulcers of the Infide of the Mouth threaten to gangrene.

### LUES VENEREA.

VENEREAL Diforders are always caused by Contact, communicated by the infected to, even, the foundest Bodies.

This Contagion produces many different Diforders, which, according to the different Parts of the Body where it fixes, go under different Denominations.

Small Ulcers appearing at the Extremity of the Penis, or on the Prepuce, are called venereal Shancres: If the nervous Papillæ of the genital Parts form little Elevations like Warts, they are called Venereal Veruccæ: If the internal Superficies of the Urethra is affected, there arises a Difficulty and Pain in making Water, named a Strangury, and a Running of a yellowish, greenish, or sometimes brownish Matter; it is then called a Gonorrhæa; if Swellings in the Groins, Buboes.

When the Virus having gained the Blood, it circulates with the Humours, where-ever it stops, it produces evils of different Kinds, for Instance, Pustules, and Spots upon the Skin, that sometimes times degenerate into filthy Crusts: Ulcers in the adipose Membrane, not giving Way in the least to such Medicines as are proper in other Ulcers; which corroding the adjacent Parts, leave profound and frightful Cicatrices: These Ulcers no sooner disappear from one Place, but they quickly shew themselves again in some neighbouring Part.

The Throat most frequently, and the Roof of the Month, are corroded little by little by this Distemper: There appears in these Parts a Spot, resembling Bacon; the Voice becomes hoarse, the Action of swallowing is attended with Pain, and the Spot we spoke of gaining Ground by little and little, destroys all the fost Parts, and at length attacks the Bones of the Palare and Mouth; which grow rotten and fall in, and leave for the Rest of Life a Desormity, for which there can be no Remedy.

This Distemper, especially if it is inveterate, attacks also the Bones, and causes Swellings on them, which, if soft, are called Tophi or Gummi; if hard, Nodes or venereal Exostoses; whence ensue very bad Caries, with intolerable Pain, worse in the Night, the Warmth of the Bed increasing it; but rather easier in the Day.

and we evile of the tree Kinds, for infringer

When the Bones are corroded even to the Marrow, the Cure is extremely difficult, and though rured in Appearance, it often returns.

This Diftemper may be eafily known by the deferibed Symptoms.

The following is a safe METHOD of Treating it.

LET the Patient take, Night and Morning, one Spoonful of the Medicine No 66. drinking after it a Pint of Barley Water with a Third of Milk in it, using the same for his common Drink: If Milk is difficult to get, the Decocion No 67. may be substituted in its Room.

This Medicine gives no Manner of Trouble to the Patients; to fome it procures fome light Stools, but this feldom; in others it works by Urine and Sweat. Its Use may be continued with the greatest Safety till all the Symptoms of the Difease have absolutely disappeared.

If the Weather is mild and temperate, the Patient may go out; but in cold or damp Weather, it is better he should keep his Room. If the Medicine feems to act too flow in robust Habits, or when the Distemper is inveterate, the Dose may be augmented to a Spoonful and a Half: And if in some Days the Symptoms do not diminish, two Spoonfuls may be given every Night and Morning, in all sour Spoonfuls a-Day.

The Time the Patient is to continue the Use of this Medicine, cannot be exactly limited; Often, if the Distemper is not very bad, the Cure is performed in three Weeks; if inveterate, it takes up more Time. But it may certainly be used a long while without the least Inconvenience.

When the Ulcers cleanse and cicatrize, when the rotten Parts of the Bone separate and fall off, and when the Tumours and nocturnal Pains diminish, the Distemper gives Way to the Remedy.

As to the Regimen of the Diet, let the Patient have Broths with Barley, Rice, or Oats, or tender Greens, his Flesh-meat lean, Milk-diet, and ripe Fruit.

Fat and smoaked or salted Meat, especially Bacon, are bad. The following Remark is necessary to be attended to. Sometimes a Salivation comes on from the Use of this Medicine, but this is but seldom, and almost only to those who have before made Use of Mercury, either internally or externally: Nevertheless, salivation not being necessary to the Cure, the Use of the Medicine No 66. is to be left off immediately on the first Signs of a Spitting coming on.

But the Decoction No 67. may be still continued.

The Signs of an approaching Salivation are as follows.

The Gums begin to swell, to grow red, to itch, and become painful, and the breath to smell ill. As soon as these Symptoms are remarked, the Use of the Remedy No 66. must, as has been before said, be suspended: But if in eight or ten Days these Symptoms disappear, and the Patient is not cured, it may again be used.

If he has a Gonorrhæa, he must drink plentifully of the Decoction No 67, to take off the Acrimony of the Urine; he may bathe the Penis thrice a-Day, for a Quarter of an Hour at a Time; in equal Quantities of Water and Milk warm. If, by Suppression of the Gonorthæa, or from any other Cause, a Testicle becomes swelled and painful, and the Scrotum red; let him be bled immediately, the Fomentation No 12, applied to the Testicle, and let him drink plentifully of the Decoction No 1, adding twenty Grains of Nitre to each Pint. After the Redness, Swelling, and Fever, which often attend swelled Testicles, are appealed, it will be proper to make Use of the Remedy No 66.

If Venereal Buboes grow hard, a Plaister of Galbanum is to be applied.

The

#### The ITCH.

I sa very troublesome Distemper to Armies, and immediately spreads by Contact, unless the affected Soldiers are separated from those who are well.

Though all the external Parts of the Body may be affected, yet the Itch most commonly shews itself first on the Hands, principally between the Fingers: First appears a Pustule, or two, sull of a Sort of clear Water, which itch extremely: When these Pustules are broke by Scratching, the Water that issues out communicates the Disorder to the neighbouring Parts. It is not easy in the Beginning to distinguish the Itch, unless one is well acquainted with this Disorder: But in its Progress the Pustules augment both in Number and Size; and when opened by Scratching, a disgustful Crust is formed, and the Evil gains the Superficies of the whole Body.

Hitherto the Itch hath its Seat between the Epidermis and the Skin; but if it continues long it makes Way through the Skin into the Membrana Adipola, where it forms small Ulcers, commonly in great Number. This Sort of Itch is the

K 2. naftieft

nastiest and worst, and at the same Time extremely contagious.

It is to be treated in the following Manner.

The Body is to be kept clean, and Linen often changed: If the Season of the Year permits, they must bathe, and in Waters impregnated with Sulphur if they can: If they cannot, Experience shews, that to bathe in Running Water will be of Service. The Shirt, Breeches, and Stockings, are to be scented with Brimstone before they are put on: But this Fumigation must be made in the open Air, lest the sulphurous Vapours should do Hurt, if taken in by Respiration.

Let the Patient take the purging Powder No 68. in the Morning fashing, and repeat it every eight Days.

On the intermediate Days, let him take, Morning, Noon, and Night, one of the Powders No 69.

Let the Parts affected be anointed every Night with the Ointment No 70.

If the Itch covers both the whole Body and all the Limbs, they must not all be anointed together at once; but you may begin by the Hands, and and Arms; continue the next Day by the Feet, Legs, and Thighs, and the third Day the Body; the fourth Day to begin again by the Hands and Arms, the fifth the Feet, &c. and so on to the entire Cure.

When the Pustules are dry, the Crusts fall, and the Ulcers disappear, and return no more, the Patient is well.

Some Spots will indeed remain on the Skin, but these Marks wear off insensibly, and in Time quite disappear.

The Patient must abstain from all Manner of falt Food during the Cure.

teams residented in a grove return who also

ou other over some bloom was direct

to the store of the second street when

THE THE WINDS

of the national residence of the first the second

K 5 WORMS

to not need the softeness of the party and the contract of the

### WORMS.

SOLDIERS are frequently troubled with Worms. Bad Food, unwholesome Water, and other Causes, engender them. Vertigæi Nausea, sudden swelling of the Lower Belly, especially after Meals, the Heartburn, Grumbling in the Bowels, and troublesome Itching of the Nose, are Indications of Worms. Some have a voracious Appetite, others lose it entirely: The Face is pale and sunk.

But all the Signs here described are not all found at the same Time in each Patient; but the more of them are discovered, the more Certainty we have of the Distemper.

But, after all, the most convincing Sign is, when the Sick void Worms, by Vomit or Stool.

All the Cure confifts in expelling them out of the Body, but this is not easy; for Worms are found to stick, as it were, to the Intestines, since otherwise they would come away with the Excrements. It will therefore be proper, in order to get the better of them, for a couple of Days to give the Patient somewhat, that by its ill Smell may, in some Manner, infect the Intestines; after which, to give a rough Purge.

Let the Patient take five Grains of Asa Fœtida in Pills every three Hours, for two Days.

The third Day let him have the Purging Powder No 71. in the Morning fasting, taking after it a light Broth, to be continued from Time to Time, till the Medicine has done working.

If, after this Process, the Symptoms do not disappear, in eight Days Time the whole must be repeated.

RECEIPES

wood iss.

003

towal and the first of the control o

All 1915, to mind on him which has will be

MATERIAL SERVICES



### RECIPES

nevel solder Jendon Deado instell

Referred to in the foregoing

or a grown When their give world track the

well in the service of the first of the contract

# TREATISE.



TAKE of the Species for the Pectoral Decoction Three Ounces; boil it in a fufficient Quantity of Water for Half an Hour, to strain Three Pounds.

The West and State of the State

Take, of the Mass for Pil. Cynogloss. Eight Grains. Make two Pills for a Dose.

Take, of the Species for the Emollient Decoction Six Ounces; boil it in sufficient Water to the Thickness of a Cataplasm\*, adding toward the End One Ounce of Mustard Seed bruised, for a Poultice.

4

Take, Flowers of Elder One Ounce; let it justboil up in a sufficient Quantity of Water in a close Vessel; then digest warm for Half an Hour; strain Two Pounds; to which add an Quace and Half of Rob of Elder, and Forty Grains pure-Nitre; Mix them together.

5.

Take, Flowers of Elder, and red Roses, of each Half an Ounce, pure Nitre One Drachm; mix them together. Infuse a Pugilt of this inwarm Water for Tea.

6.

or mostly of Water for Helf en Hopen to

Linear Three Pounds

Take, Leaves of Senna Six Drachms; Water Scurvy-grafs Two Drachms, Agaric One Drachm, and

\* A Poultice: Od a rol ell'I over salets ; sais O

+ The Eighth Part of an Handful.

and Tamarinds Half an Ounce; boil them for a Quarter of an Hour, in Water enough; strain Two Ounces, add Sirop of Rhubarb Half an Ounce. Mix them for a Draught to be taken at once.

#### 7.

Survey of the wife the remain on Owing to moth its

Take, of the Species for Emollient Decoction Four Ounces; boil it for Half an Hour in Water enough to strain Three Pounds.

#### 8.

and long the the Cincillate

Take, of the Residue after the Straining of the last, at Will, add Meal of Lintseed, Oil of Lintseed, of each Two Ounces. Mix for a Cataplasm.

#### 9

Take, Flowers of Red Roses, Two Pugils, Agrimony, One Handful; mix them together. Infuse like Tea for a Gargle. Add a little Honey.

### TO:

Take, Honey of Roses Half an Ounce; Spirit of Sea-Salt, Twenty Drops: To be mixed.

s ist made the treatment of their sa

Take, Species for the Emoffient Decoction Two Ounces; boil it in Water sufficient, for Half an Hour; strain One Pound; add Simple Oxymel Two Ounces, and pure Nitre One Drachm. Mix them for a Glyster.

### emill of most a likeling in the solid by a second

ing the Special state of the December of

Take, Species for the Emollient Decoction Three Ounces; boil it an Hour in Water fufficient; strain Four Pounds; dissolve Venice Soap Two Ounces. Mix them for a Fomentation.

#### 13.

Take, pure Nitre a Drachm and Half; Crab's Claws Two Drachms; Sirop of Wild Poppies Two Ounces; Barley-Water Ten Ounces. Mix them together.

### elenge week where a sile is to self-order

All the state of the last of the Walk of the Contract of the C

Take, Oil of Sweet Almonds, or instead thereof, best Olive Oil Two Ounces; One Yolk of an
Egg; to them well beat up, add pure Honey
One Ounce. Mix them for a Linctus.

Assert I vest Doug To a mine

Take, Kermes Mineral. Three Grains; Crab's Claws Twenty Grains. To be mixed.

16.

Take, Paul's Betony, Agrimony, Ground Ivy, Golden Rod, of each equal Parts. Infuse in warm Water for Tea.

17

Take, Cucumber Seed Half an Ounce; Eight Sweet Almonds blanched, and Two Bitter ditto. to them pounded, add One Pint of Barley Water; to be strained for an Emulsion.

18.

Take, Myrrh Fifteen Grains, and Crab's Clawa Half a Drachm. Mix them for a Powder.

19.

Take, of the Mass for Pills of Storax Six Grains, Make two Pills.

Take, Balfam Capivi Half a Drachm, and the Yolk of one Egg; to them, well rubbed in a Glass Mortar, add Pure Honey One Ounce. Mix them.

21.

Take, Coltsfoot, Scabious, Tops of St. John's Wort, of each one Handful; Powdered Liquorice Two Ounces; mix them together. Infuse for Tea.

22.

Take, Pure Nitre One Drachm; Crab's Claws Two Drachms; Sirop of Marshmallows One Ounce; and Barley-Water Ten Ounces. Mix. them.

23.

Take, Rasping of Sassafras Two Ounces; of the Three Saunders, of each Two Drachms; Powder of Liquorice One Ounce. Mix them. Insuse for Tea.

Take, Liquid Laudanum Fifteen Drops; Sirop of Diacodium Half an Ounce; and Barley Water One Ounce. Mix them for a Draught.

25.

Take, dry Camomile Flowers Two Ounces; Cochineal Twenty Grains: Boil them in Three Pints of Water to Two; in the strained Liquor, dissolve Salt of Wormwood Two Drachms.

26.

Take, Tartar Emetic in Powder. Five Grains for one Dose.

27.

Take, Root of Ipecacuanha in Powder. Half a Drachm for one Dofe.

28.

Take, of the Compound Powder of Scammony Thirty-five Grains.

Take, Sal Polychrest Two Drachms; Tartar of Vitriol One Drachm; Solutive Sirop of Roses Two Ounces; Barley - Water Half a Pint; Distilled Water of Orange Peel Two Ounces. Mix them.

30.

Take, finely powdered Jesuits Bark One Ounce; to be equally divided into twelve Papers.

31.

Take, clarified Honey Three Pounds; best Wine Vinegar One Pint. Mix them.

32.

Take, Cream of Tartar Forty Grains; Sal Polychrest Twenty Grains; Mix them. Several of these Doses to be given as Occasion may require.

33.

Take, Electuary of Bay Berries, Conserve of Wormwood, of each One Ounce. Mix them.

Take, Pill. Rusii Thirty Grains. Make seven

#### 35

Take, Oxymel of Squills Two Ounces; Sal. Polychrest Two Drachms; vitriolated Tartar One Drachm; common Water Eight Ounces; Spiritous Mint Water Half an Ounce. Mix them.

#### 36.

Take, Sal. Polychrest. Two Drachms; Tartar of Vitriol One Drachm; Electuary of Bay Berries Three Ounces; Solutive Sirop of Roses enough to make an Electuary.

#### 37.

Take, Dog-Grass Half a Pound; Dandelion, with Leaves and all, Four Ounces. Cut and bruise, and boil them in a sufficient Quantity of common Water, (or Whey if conveniently to be had, for Half an Hour; strain by strongly presenting Two Pints. Add pure Honey Three Ounces. Mix them.

Take, Sal Polychrest Two Drachms; Tartar of Vitriol One Drachm; Solutive Sirop of Roses Two Ounces; Barley - Water Half a Pint; Distilled Water of Orange Peel Two Ounces. Mix them.

30.

Take, finely powdered Jesuits Bark One Ounce; to be equally divided into twelve Papers.

31.

Take, clarified Honey Three Pounds; best Wine Vinegar One Pint. Mix them.

32.

Take, Cream of Tartar Forty Grains; Sal Polychrest Twenty Grains; Mix them. Several of these Doses to be given as Occasion may require.

33.

Take, Electuary of Bay Berries, Conferve of Wormwood, of each One Ounce. Mix them.

Take, Pill. Rufii Thirty Grains. Make seven

### 135. Hall at the same end

en linguage, cristica declara

Take, Oxymel of Squills Two Ounces; Sal. Polychrest Two Drachms; vitriolated Tartar One Drachm; common Water Eight Ounces; Spiritous Mint Water Half an Ounce. Mix them.

#### 36.

Take, Sal. Polychrest. Two Drachms; Tartar of Vitriol One Drachm; Electuary of Bay Berries Three Ounces; Solutive Sirop of Roses enough to make an Electuary.

#### 37

Take, Dog-Grass Half a Pound; Dandelion, with Leaves and all, Four Ounces. Cut and bruise, and boil them in a sufficient Quantity of common Water, (or Whey if conveniently to be had, for Half an Hour; strain by strongly presenting Two Pints. Add pure Honey Three Ounces. Mix them.

Take, Tops of common Wormwood Two Ounces; Roots of Calamus Aromaticus, Gentian, Imperatory, of each One Ounce; Bay Berries One Ounce and an Half; Juniper Berries Three Ounces; wild Carrot Seed One Ounce. Cut, bruise, and mix them. Insuse warm in a close Vessel, in, either good Wine or Mead, Eight Pints, for twenty-four Hours.

39.

Take, fresh Squills Half an Ounce. Insuse them in Two Pints of good Wine.

40.

Take, Camphire One Drachm; dissolve, by rubbing it with Oil of Sweet Almonds, One Ounce, in a Mortar.

41.

Take, Distilled Oil of Aniseed Four Drops, White Sugar Forty Grains; powdered Rhubarb Fisteen Grains, Mix for a Powder.

Take, Distilled Water of Mint 8 Ounces; Spiritous Water of Mint Half an Ounce. Mix them.

#### 43.

Take, simple Distilled Cinnamon Water One Ounce; Barley-Water Half a Pint; pure Opium Three Grains; Crab's Claws One Drachm and an Half; and, Sirop of White Poppies Half an Ounce. Mix them.

#### 44.

Take, best Rhubarb One Drachm; Citrine Myrabalans Half a Drachm. Mix for a Powder.

#### 45.

Take, Venice Treacle One Drachm for a Bolus.

#### 46.

Take, Ipecacuanha Thirty Grains, in a Powder.

#### 47.

Take, crude Opium One Grain, in a Pill.

Take, Waxed Glass of Antimony Eight Grains, in Powder.

49.

Take, Bole Armoniac Six Drachms; Gum Arabic. One Drachm; Venice Treacle One Ounce and an Half; Syrop of White Poppies, enough to make an Electuary.

50.

Take, good Wine Half a Pint; Barley - Water One Pint and an Half; Cinnamon Water One Ounce; and White Sugar Six Drachms. Mix them.

51.

Take, pure Turpentine Two Drachms; One Yolk of an Egg; to them, well mixed, add Venice Treacle Half an Ounce; pure Milk Five Ounces. Mix them for a Glyster.

Take, Species for Emollient Decoction Two Ounces; Boil it in Water enough for Half an Hour; strain Ten Ounces; add Linseed Oil Two Ounces. Mix them for a Glyster.

#### 53.

Take, Leaves of Marshmallows two Handfuls, Roots of Ditto, One Ounce; bruised Linseed Two Drachms. Boil them for Half an Hour in Water enough to strain Three Pints; add pure Nitre One Drachm; pure Honey Three Ounces. Mix them together.

#### 54.

Take, Tamarinds Three Ounces; boil it a Quarter of an Hour in Water enough to strain Three Pints; add, pure Nitre One Drachm; Honey Two Ounces. Mix them.

Take, Meal of Mustard - Seed One Ounce; Linseed One Ounce and an Half; Beans One Ounce; common Salt Two Drachms; Vinegar enough to make a Paste, to be applied to the Soles of the Feet.

56.

Take, White Vitriol One Drachm; common Water One Ounce. Mix them together.

57-

Take, Agaric of the Oak, powder it.

58.

Take, Venice Treacle One Drachm; Salt of Hartshorn Ten Grains. Mix them for a Bolus.

59.

'Take, New Milk Two Pints; generous white Wine Four Ounces; boil them for an Instant; strain the Whey from the Curd for Use.

Take, Virginia Serpentine Root, Contrayerva Root, of each Ten Grains; Bark Half a Drachm; Camphire Four Grains. Mix them for a Powder.

61.

Take, Camphire One Drachm; Dissolve, by rubbing in a Mortar, and dropping on Spirits of Wine Rectified, Twenty Drops; add, white dry Sugar Two Ounces; rub them well together, pour on White-Wine Vinegar Ten Ounces. Mix them. Keep it in a well stopt Glass Vessel for Use.

62

Take, Horse-Radish Root, fresh gathered and sliced thin, Four Ounces; Leaves of Scurvy-grass fresh gathered, Marsh-Tresoil, of each Two Handfuls; Sage, one Handful: Cut, mix, and insuse in generous Wine Six Pints, in a close Vessel, with a gentle Heat, for twenty-four Hours. Strain it for Use.

Take, Root of Sharp-pointed Dock, Polypody of the Oak, of each Half an Ounce; Chrystals of Tartar Three Drachms: Boil for Half an Hour in Three Pints of Milk; strain it; add One Ounce and an Half of Honey. Mix them.

64.

Take, Spirit of Scurvy Grass Two Ounces; Elixir of Aloes One Ounce. Mix them.

65.

Take, Spirit of Sea-falt One Drachm; Honey of Roses One Ounce and an Half; common Water Five Ounces, Let them be mixed.

66.

Take, Corrosive Sublimate Twelve Grains; rectified Malt Spirit Two Pints. Keep it in a clean Glass Vial, till the Mercury is quite dissolved.

Take, Root of Marshmallows Two Ounces; boil it in a sufficient Quantity of Water for an Hour, adding near the End, One Ounce of powdered Liquorice. Strain four Pints.

68.

Take, Scammony Fifteen Grains; fine Sugar Ten Grains; Ethiops Mineral Twenty Grains; Diaphoretic Antimony Twenty Grains. Mix them for a Powder.

69.

Take, Flower of Sulphur Thirty Grains; Ethiops Mineral. Ten Grains. Mix for Twenty-one Doses.

70.

Take, Ethiops Mineral One Ounce; Hogs Lard Three Ounces. Mix them for an Ointment.

Take, Turbith Mineral Five Grains; Root of Jalap Forty Grains; finest white Sugar Twenty Grains. Mix, and rub it to a very fine Powder in a Glass Mortar.—This Dose should be adjusted to the Age and Strength of the Patient.

Ture, Scottered I dien Granty Affrica.

1. A Granty Editor Control of the Control

Contract And


- Million the first of the last of the second to sell the second to see

สมบัญหางสารเทศ (การเกิด การเกิด การเกิด

M 2

-1-10-134-06-1733-1

A STATE OF THE STA





THE

# PRACTICE

OF

### INOCULATION

IMPARTIALLY CONSIDERED.



MANAGORNACH CONTRACTOR

H

## P.RACTIGE

INOCULATION

IMPARTIALLY CONSDERED.

E 3.5

### Practice of Inoculation

IMPARTIALLY CONSIDERED;

ITS

#### SIGNAL ADVANTAGES

FULLY PROVED;

AND

The Popular Objections against it,

In a LETTER to Sir Edward Wilmot, Bart.

By JOHN ANDREW, M. D.

TO WHICH ARE ADDED,

-

The SENTIMENTS of Dr. HUXHAM, and several of the most eminent Physicians, of both Kingdoms, relative to the said Practice,

As COMMUNICATED to the AUTHOR.

D U B L I N:
Printed by JOHN EXSHAW.

M,DCC,LXV.

Entropy of Armony ALL STATES AND A STATE OF THE STATES The state of the s There of the state of the same and TO THE OTHER POPULATIONS armore than the family to the second William In the art of the contract of the 20 40 101 THE THE HOLD MAN AND AND THE WALL WILLIAM M2.1,20.0,14



# PREFACE.

of the Proninces from a first was and



thanks a control wants aftern the stranger

\*\* HO' Inoculation was antiently practif
BA T de ed on various Kinds of Plants, with

A the Defign of meliorating their Fruit,

KR R as is evident from Virgil, Columella,

Sc. no Inflance occurs in which that Word was

used to express any Operation on Human Bodies,

analogous to the modern Method of communicating
the Small-Pox.

It is, indeed, a Matter of great Doubt, whether the Small-Pox was at all known to the Antients; for the late learned Dr. Priend Jays, the Arabian Physicians of the Seventh Century, are the first Perfors that have given any clear Description of it.

was the following the Tener surranged by chine-

From that Time, we know it has continued to make dreadful Devastations in many Parts of the World, the most learned of the Faculty having not been able to discover any Method for destroying, or even alleviating the Malignity of this most destructive Distemper.

Inoculation, like many other great and useful Discoveries, owes its Origin to the Vulgar and Illiterate; and all we know of it is, that during two or three Centuries it has been practifed in one or more of the Provinces subject to the Turks, or, as some affirm, in Georgia. But Mr. James Porter, our late Ambassador at Constantinople, assures us, that be was informed by a credible Missionary in Georgia, That he could never find any Traces of it there; and the many affirm, that it was first introduced into Circassia, with a View of preserving the Beauty of such Ladies as were destined for the Grand Signior's Seraglio, yet no authentic Evidences of this appear.

The most probable Account seems to be that, which is given by Pylarini, \* and adopted by Mr. Porter, viz. That the first who practised Inoculation was a Woman of the Morea, and that her Successor was of Bosnia. The Turks, infatuated by chimerical

<sup>\*</sup> This Person sent from Constantinople, in the Beginning of the 18th Century, to the Royal Society in London, the first authentic Account which we have of Inoculation.

rical Notions of Predestination, could never be prevailed on to submit to this Practice; but those of the Greek Church readily embraced it, and thereby it came to the Knowledge of an English Lady, \* distinguished by ber Sense, and Sprightliness of Parts, who accompanied her Husband, the British Ambassador, to Constantinople; from whence, in 1716, She fent the following Account to England, in a Letter to Mrs. S- C-, " Apropos of Diffempers, I am going to tell you a Thing, that would make you wish yourself here. The Small-Pox, so fatal and so general among st us, is here entirely harmless, by the Invention of ENGRAFTING, which is the Term they give it. There is a Set of old Women, who make it their Bufiness to perform the Operation, every Autumn, in the Month of September, when the great Heat is abated. People fend to one another to know if any of their Family has a Mind to bave the Small-Pox : They make Parties for this Purpose, and when they are met (commonly fifteen or fixteen together) the old Woman comes with a Nut-

\* Was Merit to be estimated, rather by the Nature of Things, than vulgar Opinion, this British Lady [Lady MARY WORTLEY MONATAGUE] who sirst introduced this Practice into England (by which so many Thousands of Lives have been saved) certainly deserved, and perhaps in a more grateful Age, would have had a Statue erected to her Honour, preferably to any of the illustrious Destroyers of Mankind.

Institutes of Health, P. 98.

Nut-shell full of the Matter, of the best Sort of Small-Pox, and asks what Veins you please to have opened. She immediately rips open that you offer to her, with a large Needle, (which gives you no more Pain than a common Scratch) and puts into the Vein as much Matter as can lie upon the Head of ber Needle, and after that, binds up the little Wound with a hollow Bit of Shell, and in this Manner opens four or five Veins. The Children or young Patients play together all the rest of the Day, and are in perfect Health to the eighth; then the Fever begins to seize them, and they keep their Beds two Days, very seldom three; they have very rarely above twenty or thirty in their Faces, which newer mark, and in eight Days Time they are as well as before their Illness. Where they are wounded, there remain running Sores during the Distemper, which I don't doubt is a great Relief to it. Every Year Thousands undergo this Operation, and the French Ambassador Says pleasantly, that they take the Small-Pox here by Way of Diversion, as they take the Waters in other Countries. There is no Example of any one that has died in it; and you may believe I am well satisfied of the Safety of this Experiment, since I intend to try it on my dear little Son. I am Patriot enough to bring this ufeful Invention into Fashion in England; and I should not fail to write to our Doctors very particularly about it, if I knew any of them, that I thought had Virtue enough to destroy such a considerable Branch of their Revenue, for the Good of Mankind. But that

shat Distemper is too beneficial to them, not to expose to all their Resentment, the hardy Wight that should undertake to put an End to it. Perhaps if I live to return, I may, however, have Courage to wat with them.

Stantage Correspond

In the Year 1719, Inoculation was introduced into France; but the' it was practifed there fo long fince, the' fuch a celebrated Poet as Monf. de Voltaire, and such an eminent Philosopher as Mons. de la Condamine, bave exerted their utmost Abilities in order to establish it; and the' the Children of the Duke of Orleans were fuccefsfully inoculated at Paris, feweral Years ago; yet fill it is more fparingly practifed in France, than in any other Nation in Europe, Spain only excepted: And even during the last Year, the Faculty of Physic, and the Faculty of Divinity, were employed in examining whether Inoculation ought to be tolerated or probibited. Indeed, it is faid, that it was suspended by the Parliament of Paris, and the Confideration of it referred to a Committee of Divines, and Physicians; and that swelve of the latter met, fix of whom being of one Side, and fix of the other, dijudicatio ad aliud demum tempus dilata fuit. In order to end this Difpute, I would only ask them, Whether it is lawful to fave Life, or to destroy it? That in a Country where (to use an Expression of Boerhaave) Theologorum ignorantia dominatur, Superstition Sould prevail over found rational Judgment, is no Way Jurprizing.

surprising, but that Physicians should be capable of questioning, whether the most falutary Practice that ever was discovered, should be adopted, or rejected, almost surpasses Belief. - This is the more astonishing, seeing this same Faculty of Physic (as I have been informed) applied to Dr. Huxham, and same other eminent Physicians in England, for their Sentiments of Inoculation, and received from them. the most favourable Account.

Notwithstanding Lady Montague fent such a lively Description of the Advantages of Inoculation, in the Year 1716, no one bad the Courage to put it in Practice in England, till ber Ladysbip returned in 1721, when the first Experiment was made on ber own Daughter, who having a favourable di-Riner Sort of Small-Pox, laid the Foundation of Inoculation in England. After which, in the same Year, Dr. Keith's Son, five condemned Criminals in Newgate, and three other Persons, were succefsfully inoculated, by Mr. Maitland, the Surgeon that had attended the Ambassador in his Journey to Conftantinople.

Few of the English Physicians approved of it at first. After some fort Time it met with the Countenance of Dr. Mead, and Sir Hans Stoane; but above all, the Princefs of Wales, of a Genius Superior to that of the rest of ber Sex, soon discovering obe Propriety and Reasonableness of this Practice,

gave the greatest Encouragement to it, by inocirlating her own Family; by which Means her Royal Highness preserved not the Lives only, but the Beauty of some of the most amiable Princesses in Europe.

Such an Example would probably have been followed by Perfons of all Ranks, if an unfortunate Event had not happened at this Juncture. — A Son of the Earl of Sundetland, who was inoculated from after the Royal Family, died; and the his Death was proved to have been owing to an Hydrocephalus, or Collection of Water in his Brain, it helped to furnish Objections to fuch as were previously Enemies to the Practice, but notwithstanding it fill met with Encouragement, the much less than was expected; the whole Number of the Inoculated in England during the three first Years amounting only to 477.

Lady Montague, was much decrived, in thinking, she should be obliged to war with the Faculty, on Account of her Endeavours to introduce Inoculation. The principal Auxiliaries † she met with N 2

<sup>\*</sup> The Royal Family were inoculated in the Beginning of the Year 1722.

<sup>+</sup> As the Faculty have taken fuch Poins to offablish a Practice that is manifefully injurious to their

were the Physicians themselves, among whom no one distinguished bimself so eminently as the celebrated Dr. Jurin, who being possessed of the greatest Share of Claffical and Philosophical Learning, as well as the most profound Medical Skill, employed bis great Abilities in making the most accurate Obfervations on the Effects of this Practice. He by procuring authentic Narratives of the Success with which it was attended, and comparing them with Lifts of those who died of the Small-Pox in the Natural Way, gave the most convincing Proofs of the Preference that ought to be given to Inoculation. His Writings carried with them irrefistable Conviction: They enlightened the Understanding of the Ignorant, confuted the Errors of the Mistaken, and in great Measure, belped to remove the Fears, and Doubts of the Timorous and Superstitious. He publifbed Annual Accounts of the Success of Inoculation. during several Years; but from the Time of bis discontinuing bis Narratives, till the Year 1746. this most excellent Practice feemed to decline, in London. In that Year it regained new Life, and Vigour, by the Establishment of the Hospital for Inoculation

### NOTE.

own Interest, it is hoped the World will be convinced of their Sincerity, and confess, that their Representations of the Benefit of it to Mankind are such as may justly be depended on. ne

t-

eft

ms.

ed

b-0-

3

3

e

r.

oculation; to which Persons of the greatest Rank, as well as many of inferior Order, soon became Contributors, at the Head of whom is an illustrious Personage, who embraces all Opportunities of encouraging, and protecting, every laudable Scheme for the Benefit of Mankind.

Within the Period just mentioned, I began the Practice of Inoculation in the West of England, where I found the Prejudices against it so great that I was obliged to practise as it were in the Dark, wisting my Patients only by Night, and doing Good by Stealth, as will be made appear by a Letter from Dr. Alured Clarke, our late worthy Dean, who founded the Devon and Exeter Hospital, in the Year 1741 I could wish such a laudable Example might be imitated; and that an Hospital was established here for Inoculation. Such an Undertaking could not sail of being encouraged by the principal Inhabitants of this great and opulent City and County, many of whom have experienced the Benefit of Inoculation in their wn Families.

N<sub>3</sub>

THE

See 30 V to 1 A 1.



# to the regular to B H to C to the rest of the manager which

redictioned to properly bearing

# PRACTICE

O F

# INOCULATION,

IMPARTIALLY CONSIDERED.

# 

To Sir EDWARD WILMOT, Bart.

A se me the first Notion of Inoculation, and as my Method is principally founded on your Precepts, you will give me Leave to acquaint you with the Success that has attended it.

stool plater office or erect of such alight

2

The Difficulties I met with in the Beginning of my Practice, may partly be gueffed at, by the following Letter from Dr. Clarke, Dean of Exeter, who defired me to inoculate one of his Servants. Though I had from my first Entrance on the Study of Physic, and my early Instructions from you, a very high Opinion of Inoculation, I was averse to the putting it in Practice in Exon, without a positive Order under the Dean's own Hand, which occasioned the following Letter.

#### To Dr. ANDREW.

Dear Sir,

Winchester, Nov. 19, 1741.

WITH Regard to Margaret, I do here under my Hand, clear the Doctor and Apothecary of all intentional, or actual Guilt, of any Thing that may happen to her, in the Course of their lawful Employments. When I left Exeter, she was very desirous of being inoculated. I hope nothing has happened that can intimidate her. A Doctor tame here tother Day, and produced a List of sixteen hundred Persons, who were inoculated between Farnham and Chichester, and the intermediate Country, of which I believe there were not six Milicarriages. In short, if you think

Leave to acquaint from with the Success

<sup>\*</sup> The Dean was very fearful of exceeding the Truth, that he appears to have fallen greatly thort

her a proper Subject, and she is really willing to submit to the Operation—I earnestly defire it; and doubt not of God's Blessing on so reasonable an Assempt. But it is with this sine qua non, that no-body, not even her Fellow-servant, may know it is by Inoculation, till the Prejudices against it are a little more worn off.

I am, Ge.

A. CLARKE.

## NOTE.

of it, as is evident by the following Note, printed in the Bishop of Worcester's Sermon, which was preached before the Governors of the Hospital for Inoculation, March 1752, and evidently refers to the same Account as that given by the Dean, in his Letter to me. " The learned Dr. Langrish, of Winchester, gives the following Account. - About ten Years ago the Small-Pox raged fo furiously in Winchester, and the neighbouring Towns, that great Numbers were induced to be inoculated. In Portsmouth, Chichester, Guilford, Petersfield, and Winchefter, we reckon near Two Thousand that were inoculated at that Time, and (except two Women, who were both with Child, and inoculated contrary to the Advice of their Phylicians) I don't remember to have heard of one bad Accident that befell any of them."

4

This Margaret was the Dean's Cook, aged 41, very corpulent, and, in Appearance, an improper Subject for Inoculation; however, the went thro' the Diftemper without any bad Symptom.

This, and some more Instances of the like Kind, encouraged me to be bolder, and no longer to practise (even in Exon) in the secret Manner enjoined by Dean Clarke. I think proper to mention one other, before I proceed to give an Account of my Method, and its Success.

Mr. Cholwich, a Gentleman of great Worth, and confiderable Fortune applied to me for Directions concerning the Inoculation of his Children. The Success of which appears by the following Letter,

To JOHN CHOLWICH, Efq; in Exeter.

Dear Sir.

Oldfton, Dec. 1. 1741.

A S you were so kind as to give me your Advice, and Approbation, on the Inoculation of my Children, I do not doubt but it will be agreeable to you, to hear of our Success. November the 10th, after about three Weeks Preparation, the Operation was performed on all three.

the Advice of their Paris Third cor's consumer

My Son continued very well till the feventhe Day after the Operation, when he complained of a Pain in his Head, which was his only Complaint, and that not violent, till the Eruption, which was on the ninth Day.

My Daughters grew sick the Day after their Brother, and for two Days were delirious at Times; but from the Time of the Eruption, which was on the tenth, all thro' the Distemper were very calm and well.

The Number of Pustules in the Face, did not exceed Sixty in either of them, neither had they many in their Bodies. They turned the seventh Day in all, and without the least bad Symptom of any kind.

The Diforder proving so exceedingly favourable, I had not any Occasion to confine them to their Beds, after the Eruption. They eat, drank, and slept very well during the whole Distemper, and had scarce any Occasion for a composing Draught. The tenth Day from the Eruption, they were all so well as to be purged; and are now, I bless God, as well as I could with.

Upon the whole, I am convinced, the Method of Inoculation is a very fafe and ufeful Practice; and as such, I cannot help withing it may meet with a more general Approbation, especially as

the Small-Pox hath been so fatal of late in this County. —— If you think the Example I have set, will incline others to imitate it, you may make this Account as public as you think proper. I am, with due Compliments to your whole Family,

Your most affectionate Kinsman, and

Most obliged bumble Servant,

#### WILLIAM CHOLWICH.

Though Inoculation has been introduced into Exeter, and the County of Devon, more than twenty-three Years, as appears by the above Letters, it is still very sparingly practised; and the Prejudices against it being not yet worn off, I am induced to publish this short Treatise.

The Inoculated Small-Pox, seems to be attended with less Danger, than the Natural, because the Infection is, or always may be communicated by mild Pus, through slight superficial Incisions, to sound healthful Bodies, properly disposed for its Reception, in the best Season of the Year, and at a Time of Life that is apprehensive of no Danger; and because a proper Regimen is observed during several Days previous to the Disorder: Whereas the Natural may seize weak or distempered Persons, at an advanced Age, and in a Season of the Year, either violently Hot, or intensely Cold; be communicated from the most virulent Sort, by extremely

7

tremely acrid fetid Effluvia, taken into the Lungs, and Stomach,\* or foon after Intemperance, or Excels of any Kind, hard Labour, or fatiguing Journies.—Is it, then, a trivial Concern, that all the above Advantages may be obtained, and the unhappy Circumstances effectually prevented, by Inoculation?

t,

Whenever the Person to be inoculated, is of a full sanguine Habit, I always order Blood to be drawn, and generally give three or sour Doses of Purging Physic, with a sew Grains of Calomel, at proper Intervals, and an alterative Medicine, with some Preparation of Antimony every Night, on the Days that no Purging Physic is taken. This has been my general Practice, and, as such, I venture to publish it; the I am sensible it may be liable to many Objections.

Dr. Wall, of Worcester, has favoured me with the following Strictures on it, in a Letter, dated Dec. 17, 1764.—" As to the Use of Mercury, and Antimony, in the Preparation of Persons to be inoculated, I confess I have some little Doubts. I agree, that they may be extremely useful as Evacuants, to carry off stagnant Bile, or unload the Viscera

<sup>\*</sup> From this Cause proceeds the Sore Throats, Difficulty of Breathing, and violent Vomitings, so frequent, and so fatal, in the Natural Small-Pox.

Viscera of any putrid Saburra, which they effect better than any other Articles in the Materia Medica. But I should hesitate a little as to their Use, if given only as Alteratives, especially with regard to Mercurials, which are known to break down the Red Globules, and dissolve the Crass of the Blood. — Indeed to speak plainly, I don't think there is one Case in ten that requires any further Preparation, than some gentle Evacuations, and a little Circumspection in regard to Dict and Regimen; and in those Persons, where a nicer Preparation is requisite, it seems impossible to six any one Method, which shall suit the different Constitutions."

To these Observations of my worthy Friend, I will reply in the Words of Dr. Huxbam, who gave the following Answer to some Queries I sent him, concerning the Use of Mercurials, and Antimonials, as Preparatives for Inoculation.

Plynoutb,

Plymouth, Jan. 23, 1765.

" So long ago as 1724, I suggested, \* that Mercurials, as well prepared Calomel, or the like. might be of Use in the Small-Pox. I seldom fail of giving a Mercurial Purge, + or two, (but with no heating Drastic Cathartic) previously to the Inoculation of a Person for the Small-Pox. Sometimes also I give my Antimoniated Æthiops, especially when I suspect a Verminous Seminium, foul obstructed Viscera, or Glands .- The Use of Mercurials, and Antimonials, will more fully appear, as preparatory to the Inoculation of the Small-Pox, by what the ingenious Dr. Benjamin Gale, of Connecticut, in New-England, hath communicated to me, in his Differtation on the Inoculation of the Small-Pox in America; in which he fays, before the Use of Mercury, and Antimony, in preparing Persons for Inoculation, one of 100 of the Inoculated died, but fince only one of 800."

But notwithstanding Dr. Huxbam has furnished me with so good an Argument in support of my Method of Preparation, I must not conceal what he surther observes "When he has healthy O 2 young

<sup>\*</sup> Phitosoph. Transact. No 390.

<sup>†</sup> Sir Edward Wilmot's Practice corresponds, in this Respect, with Dr. Huxbam's, especially in regard to Children.

young Persons to Inoculate, he seldom gives more than a sew gentle cooling Purgatives, enjoining them to a light easy Diet, Abstinence from Flesh, or Fish, for a sew Days, and carefully guarding them against violent Exercise, and taking Cold."

Thus does he feem to agree with Dr. Wall; and the little Necessity there is for Preparation, is confirmed by the Account I received lately from Dr. Swan of Newcastle, who informs me, that in his Neighbourhood 70 or 80 Persons were Inoculated without the least Preparation, and all recovered of the Small-Pox.

But with great Deference to the Opinion of fo good Judges, as it is impossible always to be certain, whether there may not be a Verminous Seminium, as Dr. Huxham calls it, I think it not improper to give a Mercurial Purge or two, even to young Persons, who in all Appearance are quite healthy. I have at this Time (Feb. 16, 1765) under my Care a young Man, who told me, when I began the Preparation, he had no Complaint of any Kind, neither could I discover the least Symptom of any, yet by taking two Mercurial Purges, he discharged several Worms. And, it being very necessary to take particular Care, that Children be free from Worms, at the Time of Inoculation, besides the Medicines recommended above, I fometimes give them Pink Root, or Bear's Foot. \*

<sup>\*</sup> Our great Botanist, Dr. Hill, says, he is very

This last is certainly one of the best known Vermisuge Remedies, though on Account of the bad Effects that sometimes attend it, not sufficiently made Use of.

During the Preparation, I allow my Patients to eat Flesh only at Dinner, every other Day, and then nothing but Veal, Chicken, and the like, and the intermediate Days, Puddings, Fruit Pies, Rice Milk, &c. and for Breakfast and Supper, whatever they choose, provided they eat no Flesh. A Glass or two of Wine, or Wine and Water, grown Persons may be allowed at Dinner.

The above is all the Preparation that I have generally found necessary, but it may not be amiss to mention, that Dr. James says, his Powder, given in small Doses, as an Alterative, is an excellent Preparative for Inoculation.—As being a Preparation

# NOTE.

confident that all the Mischief, which is ascribed to this Plant, has been in Consequence of giving one Species for another:

One being absolutely Sase, and Mild, in it's Effects; the other a Rough and Violent Medicine, and frequently deleterious. The true Sort, is the third Article, under Helleborus, in Linnaus's Species Plantarum, a low Plant. The deleterious Plant, is the fourth Article, and is two or three Feet high.

tion of Antimony, I cannot disapprove of it, and I imagine it may be of Service in a viscid State of Blood.

The Preparation being finished, the Operator must make an Incision in each Arm, with a Lances, just deep enough to wound, but not to penetrate the Skin, about Half an Inch long, into which is put a small Thread, impregnated with the variolous Matter, and bound on pretty tight, during sixteen or eighteen Hours.

The Thread, by being drawn with a Needle through three or four of the Pustules, when fully maturated, will be sufficiently impregnated. The Incisions are to be drest every Day with yellow Basilicon, or (if there should be any sungous Flesh) with dry Lint. After the Operation, the same Regularity with regard to Diet, as already mentioned, must be continued till the fixth Day, and then I repeat the same Dose of Purging Physic, recommended for the Preparation, but without the Calomel.

After the fixth Day, I forbid all Flesh, and Wine, till the Eruption is completed, and do not permit my Patients to go off the Floor where they are, lest by any Exercise they may heat their Blood, and thereby increase the Fever, which usually begins about the End of the seventh, or Beginning of the eighth Day, after the Operation, and sometimes

times with as much Violence, as in the natural Small-Pox. From this Time I confine my Patients to their Beds, till the Eruption.

When the Fever comes on, my Method varies, as the Case may require, it being sometimes necessary to bleed, purge, give Emetics, Glysters, &c. but If the Symptoms are mild, few Medicines will be required.

As foon as the Pustules appear, (the Time for which is usually at the End of the ninth, or Beginning of the Tenth Day from the Operation) my Patients are frequently as free from all Complaints, as in the most perfect Health; and after the eleventh Day I permit them, whether they are grown Persons or Children, to rife every Day, and stay out of Bed several Hours; to take weak Chicken, or Veal Broth, every Day for Dinner; and grown Persons, if they earnestly defire it, may be allowed to eat a Bit of Chicken, Plaice, or Whiting; and to drink a Glass or two of Wine and Water; and, from that Time, I give but sew Medicines, if the Disorder proves savourable.

The Small-Pox by Inoculation, generally turn on the feventeenth Day after the Operation, and then the Wounds, made by the Incisions, begin to discharge plentifully; to which Discharge, many, how justly I cannot say, attribute the Favourable-ness of the Symptoms on the Turn. About the twentieth,

twentieth, or twenty-first Day, I repeat the Purge, and then permit my Patients to return to their usual Way of Living, strictly enjoining them to take four or five Doses of purging Physic, at proper Intervals.

Some grown Persons, being not capable of getting the better of their Fears, in case they are restless, I give them at Night twelve or fifteen Drops of Liquid Laudanum, in a Saline Draught; and to Children a Drachm, or two of White Poppy Sirop. If my Patients should be three Days without a Stool, (which feldom happens) whether they are grown Persons, or Children, I order for them an Emollient Glyster.

During the Course of Twenty-three Years, I have inoculated more than three hundred Persons. in the above Method, not one of whom has mifcarried; and in my whole Practice I have only loft one, whose Case Jeing frequently mentioned to the Prejudice of Inoculation, it may not be improper to give some Account of it here.

This Person was a promising Youth, of nine Years of Age. He had been prepared at his Father's House, about thirty Miles distant, in the usual Manner (except the Omission of Calomel;) and being brought to Exèter, the Operation was performed September 27th, 1751; from which Time till October 3d, he remained very well. In the the Evening of that Day, the Symptoms, which usually precede the Eruption of the Small-Pox, came on with some Degree of Violence, and continued, without any Remission, till the 7th in the Morning, when he brought off two large Worms; after which a Gleam of Hope appeared, but this was of a very short Duration, for the most alarming Symptoms soon returning with greater Violence, in the Evening of the same Day, he died. The perpetual Vomiting he was afflicted with (from Worms, as I suppose, in his Stomach) made it impossible to give him the least Relief; and to them I attribute his Death, for the few Eruptions that appeared, seemed to promise a sayourable Sort of Small-Pox.

Fourteen or fifteen Years ago, it was not a conflant Rule with me to give any Mercurial Medicine, during the Preparation; but fince this unfortunate Event, I have seldom, if ever, omitted it, except in very young Children, and uninterrupted Success recommends the Practice.

I cannot avoid mentioning, that as this young Gentleman refided at fo great a Distance from Exeter, I was not defired to see him during the Preparation; but since the above Accident, I have made it an invariable Rule, never to undertake the Inoculation of any Person I don't visit, and am thoroughly satisfied is properly prepared. If a Visit had been desired, I should perhaps have ordered

Bear's

Bear's Foot, Calomel, or fome other Vermifuge, which might have prevented the Train of bad Symptoms that proved fo fatal to him.

I find, among my Notes, that the next Persons
I inoculated, were four Children Inhabitants
of this City; and as their Case was very remarkable, I will give a short Account of it.

Oct. 2, 1751, A Gentlewoman applied to me to inoculate four of her Children. As they all appeared to have Worms, from their swelled Bellies, pale Countenances, Loss of Appetite, and setid Breath, I gave them at different Times, several Doses both of Calomel, and Bear's Foot; by which during the Course of their Preparation, they discharged one bundred and thirty round Worms. These Children were inoculated the 21st of the same Month, and had the Small-Pox so favourably, that the Number of Pustules in all, scarce exceeded one hundred.

Let any one reflect, what might have been the Consequence, if the Small-Pox had seized them before the Worms had been discharged, and how hazardous it is to inoculate Children, without giving them Calomel, Bear's Foot, or some other powerful Vermifuge Medicine!

The House of har his entered is used that it is a recent.

Among all that I have inoculated, not more than ten have had any confiderable Number of Puftules, only three have had the confluent Sort, and even these three had scarce any secondary Fever.

Persons who have the natural Small-Pox in a severe Manner, are generally afflicted with a fore Throat, and a Difficulty of Breathing, which, those who are inoculated, are generally either quite free from, or so slightly affected with, as to find little or no Inconvenience from them, and by this Means they are secured from the two most fatal Symptoms attending this Disorder.

I most earnestly advise all Persons to avoid the Natural Small-Pox as much as possible, which I think full as infectious as the Plague, and the worst Sort of it much more grievous to be born, and productive of worse Effects to such as survive the Distemper. I have such an Abhorrence of it, that I should live in perpetual Misery, if I thought there was the least Probability of being infected with it a second Time.

This most dreadful, and most destructive of all Distempers, is rendered so mild by Inoculation, as scarce to deserve the Name of a Distemper; tho' I hardly think the French Ambassador's Obser-

vation will be verified, That People will take the Small-Pox as they drink the Waters, for their Diversion; and yet I remember some Years since, I inoculated sour young Men in one House, who played at Cards every Day during the Disorder, except while the Eruptive Fever lasted.

The Lady who introduced Inoculation into England, says, " There is no Example of any one that has died in it :" And I have heard, it is the Opinion of feveral Persons in England, that no one dies of Inoculation. Whether this is true, or not, may be a Matter of Doubt; but I am firmly perfuaded, if Inoculation should be universally practifed, all Apprehension of Danger from the Small-Pox (if not the Diforder itself) would be annihilated in a few Years .- I have practifed it on Perfons of almost all Ages, from that of three or four Months to fifty or fixty Years, and to judge by the Success, it may be difficult which to prefer; but when my Opinion is asked, I recommend any Time between three and twelve Years of Age, and particularly advise all Females to be inoculated within that Period. With regard to the Season, I have been concerned with Inoculation every Month in the Year, but I prefer the Spring and Autumn.

Some of the Advocates for Inoculation mention it as an Advantage attending the Practice, that it has the Choice of Subjects, as if none were to undergo

dergo the Operation, but those who are healthy, strong, and of a sound Habit. These Advocates derogate from the Esteem that ought to be paid to it. I have inoculated several of an highly scorbutic, and even of a scrosulous Habit, who have had the Small-Pox as savourably as any I have been concerned with; but then I confess they have been longer in a Course of Preparation: And so far I am from choosing my Subjects, that I cannot recollect ever to have refused more than one or two Persons that desired to be inoculated, for I think it is the Duty of every Physician to inoculate all that offer, if they are in Health, except Women with Child.

Sucking Children\*, who are born of healthy Parents, feem at first View the most proper Subjects,

\* "Dr. Mackenzie says, the most favourable Period for Inoculation, seems to be that which precedes the Breeding of Teeth in Children.—Dr. Gatti says—L' Age le plus propre pour l'Inoculation est le plus tendre: But I have lately been informed by Dr. Young, of Edinburgh, that though he had inoculated a great many Infants of eight and ten Days old, he could never bring out an Eruption like the Small-Pox; and therefore, he says, the most proper Age for inoculating Children, is from two to sive Months; but he allows, it was long reckoned, that Children of three Years old were of the most proper Age for Inoculation."

jects, as fuch are free from impure Juices; but I always decline the Inoculation of them, because it is extremely difficult to give them any affistance by Medicine, in case they should stand in Need of it, and the Anxiety of the Nurse frequently spoils her Milk, by which Means a Disorder, perhaps as dangerous as the natural Small-Pox itself, may be produced, of which I have seen some Instances in my own Practice, and have been forced (in order to preserve the Life of my Patient) to procure a new Nurse in the Midst of the Disorder.

Another Argument against inoculating young Children is, that more than One-third Part of all who are born are supposed to die before they are 2 Years of Age (at least fuch is the Calculation within the Bills of Mortality) and confequently the Risk of having fome other Diforder with the Small-Pox, is greater at that Time of Life than at any other, which, perhaps, may be the Reason why Inoculation has been found fo unfuccessful in young Children; for it appears, by Dr. Kirkpatrick's Analyfis, P. 125, That of those who have been inoculated under two Years of Age, more than one in ten have died: And Mr. Scheuzer fays, that in the Year 1728, one hundred and twenty-one Perfons were inoculated, of whom three died, which is one of the most unfavourable Accounts that have been published. But then he adds, of these three, one was only eight Months old, another eleven Months, and the third a Year and half:

And by Accounts received lately from North-America, by Dr. Huxham, and by him transmitted to me, I am informed, that only five have died of three thousand who were inoculated: And the same Accounts say, that these five were all young Children.

Reflecting on the above Particulars, I recommend it to all Parents to keep their Children from Infection (if possible) till they are three Years of Age. All my own, in Number five, were inoculated between the Age of three Years and feven, and had not fifty Pustules in their Faces amongst them all. My Wife indeed, was not inoculated till she was thirty-five; but though so far advanced in Years, and though the Eruption was large, yet she went through the Disorder without the least bad Symptom. Having therefore experienced fuch fignal Advantages from Inoculation, in my own Family, I cannot any longer deny myfelf the Satisfaction of thus publicly recommending it to others and especially, as I have never known, during the thirty Years of my Practice, the Natural Small - Pox fo general, or fo fatal, as during the last fix Months.

Having, I hope, fully proved the fignal Advantages attending Inoculation, I will next endeavour to confute the most popular Objections against it. The chief Argument urged by foolifbly fond or fuperstitious Parents, against this Practice, is, that it brings a Distemper upon their Children, which they might never have, and that if any one of them should die, they should never forgive themselves, on Account of their having (as they term it) presumptuously tempted Providence.

In order to confute this Objection, I beg proper Attention may be paid to the following Remarks: -The present Bishop of Norwich, in his Sermon on Inoculation, fays, " To venture upon an Action, certainly known to be displeasing to God, as having been directly forbidden by him, is, in Scripture Language, tempting God. But before fuch an Action as this is condemned, as a Temptation of God, you must prove it to be actual Difobedience, or certainly so by direct Inference; which neither has been done yet, nor I believe ever will be done. Nor can Inoculation be called not depending upon Providence, any more than every other prudential Act, especially every other preventive Act, which human Wisdom, and Forefight, may and daily does suggest."

The present Bishop of Gloucester, in his Sermon on the same Subject, has the following Observation: — "By what strange Fate it is, I know not, says his Lordship, but so it has happened, that, at a Time

a Time when Religion has loft almost all its Influence on the Minds of the People, Superstition still keeps its hold; and this most beneficial Practice of Inoculation is regarded with Abhorrence by them, as a Kind of Impiety, a tempting God, and miftrusting his general Providence. Now the best Confutation of such monstrous Absurdities, is the Success of the Practice."

The Rev. Mr. Some fays, " 'Tis obvious that we are to depend upon the Care of Providence, only in the Way of our Duty. To talk confidently of our trufting God, while we omit the proper Means of escaping the Dangers which surround us, is not Faith, but unwarrantable Presumption."

Thus far these learned Divines -- In a Word, Providence has given us Reason to discover what is most for our Advantage, and has furnished us with a Power to execute it: To neglect, therefore, the Use of them for that Purpose, is not to trust in Providence, but to be presumptuous, and ungrateful.

I will next consider the great Risk of having the Small-Pox in the Natural Way, as I am fenfible the pleasing Imagination of being able to avoid it, induces several to neglect this most beneficial Practice. But if such Persons would reslect, that there is scarce one in a thousand that escapes this

P 3

Diftemper.

Distemper, (among those, I mean, that frequent Public Places, and converse familiarly with Mankind) and that at least one in five dies, of those who have it after they are arrived at Years of Maturity, I believe they would change their Way of Thinking.

If my Readers will attend to the Observations here subjoined, and which are taken from the most celebrated Practitioners in England, France, Italy, Holland and Germany, they will be convinced that my Account of the Danger of Insection, is no way exaggerated.

Riverius \* says, "You can scarce find one of a thousand, who once in his Life is not afflicted with this Distemper." Sennertus says, "This Disorder is so common, it is difficult to find any one that has not, at some Time or other been infected with it."——Sydenham says, "It seizes whole

\* Ex hominum millibus vix unum reperire licet qui semel saltem in vitâ hunc affectum non patiatur. Riv.—Variolæ omnibus hominibus adeo communes sunt, ut vix reperiatur, qui non aliquando iis inficiatur. Sennert.—Variolæ integras familias contagio suo adflantes, nimini parcunt. Syden.—Vix ullus alius morbus, exceptâ forsan pefilentia, adeo contagio sus est, ac variolarum. Hoffman.—On convient que presque tous les hommes ont la petite verole une fois en leur vie. M. Gatti.

whole Families, without fparing a fingle Person." -Hoffman fays, " No Diftemper, except perhaps the Plague, is fo contagious."-Monf. Gatti fays, " It is allowed, that almost all Persons have the Small-Pox once in their Life." -Dr. Mead reckons scarce one in a thoufand escapes it.

Tho' the above learned Phylicians represent the Danger of Infection to be fo great, it is not to be prefumed, they believe that not more than one in a thousand dies, without having had the Small-Pox; their Meaning is, that very few go through Life without having this Diforder; for, according to the most exact Calculations, one half of all who are born never have the Small-Pox at all, and of the other half, as one in feven dies of this Distemper, it carries off one fourteenth Part of Mankind, which is much more than can be faid of any other Distemper; and as Inoculation frees us from this dreadful Calamity, it may justly be reckoned the greatest Bleffing that ever was discovered.

If the Small-Pox is really fo contagious as the above Practitioners reprefent it, how abfurd are those who expect to avoid it? And how can Parents reconcile it to their Conscience, to let their Children run fo great a Risk of dying of the Small-Pox in the Natural Way, fince Providence has made known an almost infallible Method of securing them from all Danger in that Distemper, and thereby avoiding what may be reckoned one of the chief Scurges of Human Nature?

I repeat it, that Inoculation is almost infallible, and if its Success was more certain than it really is, Mankind might be inclined to trust entirely to their own Skill, and lose all Dependence on the Deity. Some Authors, however, represent it as quite infallible. — Dr. Timoni, who practised at Constantinople, says, in one of his Treatises on this Subject, that "not one Person of those who were inoculated died." And in another, that "during a Course of seven Years, and even when the most malignant Fevers raged, he had seen it practised on Persons of every Age, Sex, and Condition, without the least bad Accident of any Kind."

Pylarini, who practifed also at Constantinople, makes the same Observation, and in the same Words, adding, that "every one may be assured, when Inoculation is conducted in a proper Manner, and after the necessary Preparation, it promises Safety that is morally certain."

But why should we have Recourse to Foreigners for the Recommendation of a Practice, which has been so successful in our own Country? If the Superstitious and Timorous would give themselves the Trouble carefully to peruse the excellent Sermons published by several very eminent Prelates of

the Church of England, on Purpose to recommend Inoculation, I am convinced, no Scruple of Conscience, nor Apprehension of Danger, would any longer deter them from pursuing a Practice which promises such certain Benefit.

By perusing these Sermons, they will find every religious doubt resolved that can be suggested against Inoculation, and they will be convinced, that of those who are inoculated, only one in 343, and of those who are admitted into the Hospital with the Natural Small-Pox, more than one in four dies. If such is the Fate of these miserable Objects, when supplied with Food, Physic, and every Convenience suited to their unhappy Circumstances, guess the Condition of those (and many fuch there are) who are destitute of almost all the Necessaries of Life: At this Time of their greatest Distress, every Ear is deaf to their Solicitations, every Door is thut against their Petitions: -This being a true Representation of some of the Distressed, perhaps, in every Parish in the Kingdom, I think it incumbent on the Overfeers of the Poor, to take Care, that all who receive their Pay may be inoculated at proper Seafons.

A Thought just now occurs to me, which feems not unworthy of the Attention of the Government. - The Indians in North - America are greatly exasperated against the English. The Small-Pox (as we are informed by the public Papers)

now rages fo violently there, that scarce one recovers who is feized with that distemper. If the English could contrive to introduce Inoculation among them, it might be the Means not only of preserving their Lives, but that of their yielding a voluntary subjection; and this is the more needful at present, if it be true, that the French have made them believe, the English infected them with the Small-Pox on purpose to destroy them.

Of all who are inoculated, if only one in 343 dies, \* it may not be unfair to suppose, that the Person who falls a Victim, might be infected with the Natural Small-Pox, or fome other mortal Difease during the Inoculation. Monf. de la Condamine fays, that of 320 Perfons, taken at a Venture, it is probable that one will die within 40 Days. It therefore follows, that of 320 Perfons inoculated of all Ages, one is to die within that Term: If Monf. de la Condamine's Calculation is just, it behoves us to be thoroughly persuaded that no more die of Inoculation, than are absolutely neceffary to keep up a proper Reliance on Providence, and with becoming Faith and Confidence to make proper Use of a Discovery which Dr. Squire (the present Bishop of St. David's) says " is the Gift of

<sup>\*</sup> See the Bishop of Norwich's Sermon, preached before the Governors of the Inoculating Hospital last Year.

of God's Providence to Mankind."—A Discovery that future Ages will envy us, and will be aftonished at the Opposition that has been made to it. Let such as neglect to make Use of so great a Blessing, reslect, that they are counteracting the Designs of that Being, who permits this Practice to be attended with surprizing good Success.

May it, therefore, make a lasting Impression on my Readers, That if Inoculation should be universally practifed, it would be the Means of saving the Lives of many thousands, who will be devoted to certain Death every Year, in England only, by the Means of Ignorance or Prejudice; and I hope the Time is at Hand, when Parents will think they ought never to forgive themselves, if they should neglect to make a right Use of such a providential Discovery.

I hear some Parents say, they will let their Children be well prepared, and then carried to a Person who has the Small-Pox in a sayourable Manner—Nothing can be more absurd, nor inhuman than this Practice, for by it they expose their Children to a much greater Hazard, than in permitting them to be inoculated, and alike tempt Providence.

<sup>\*</sup> De la Condamine says, "In France 28,000 Persons are devoted to Death yearly, which it would be in our Power to preserve by Inoculation."

Providence, by bringing on them a Distemper that they might never have; and lest they should lull themselves in the Security of being able to procure, by this Means, a savourable Sort, I beg it may be remembered, that Preparation avails much less towards mitigating the Severity of the Natural Small-Pox, than unexperienced Persons may apprehend, the Success of Inoculation depending principally on the Manner of communicating the Insection; for which Reason, I am intirely of Dr. Heberden's Opinion, That it is better to have Inoculation performed by any Body, and in any Manner, than to suffer the Disease to come on in the Natural Way, though affished by all the Help that Art can afford.

If any should be offended with the Boldness of the Expression, Devoted to certain Death, I must acquaint them, that every Method hitherto discovered for curing the Natural Small-Pox, frustrates our Hopes; and whether we practise in the coolling Method, recommended by Sydenham; give the heating Medicines, prescribed by Moreton; or pursue the middle Tract, with our celebrated Countryman, Dr. Huxbam (which seems the most rational) or, with Friend, endeavour to purge off the secondary Fever; we shall still find, I believe, the Small-Pox will carry off near one in seven of those who have it naturally.

The Danger attending this Distemper made Sydenham declare, that the Method of curing it rests on a slippery Foundation, and verifies the Observation made by Boerhaave, that, Vulgata methodo nullus nisi sponte emergit, \* by which he means, Nature not Art performs the Cure.

I might have cited many other Testimonies to the same Purpose, it being certain that there are almost as many different Sentiments on the Treatment of this Disorder, as there are Authors who have written on it; what is approved by one being condemned by another, and the Method adopted in one Country, being rejected in another, which is a full Proof of the Difficulty at least that attends the Method of curing the Natural Small-Pox, and seems sufficient to render the Practice of Inoculation universal.

I will next endeavour to confute an Objection which has been insisted on by some of the most learned of the Faculty, even such as are great Advocates for the Practice, and therefore demands our serious Attention. Mons. Gatti says, in his Q

\* Immediately after the above Observation, he adds—Prophylaxis insitiva videtur satis certa tutaque.—Thus did this great Man Forty Years since, and almost in the Insancy of its Practice, (at least in this Part of Europe) pronounce Inoculation to be sufficiently safe and certain.

Treatise on the Prejudices to the Practice of Inoculation-(Il est certain qu'on peut communiquer avec la petite verole les maladies qui sont centagieuses, la rougeole, par example la fieure scarlatine) -" It is certain, That with the Small-Pox, contagious Diforders, fuch as the Measles and Scarlet Fever, may be communicated."-The most fatisfactory Answer that can be given to this Objection is, that I believe no Inoculator in England, where the Practice has been more general than in any other Part of Europe, ever met with an Instance of this Kind; yet I own, that in the Course of my Practice, I have frequently feen a Rash, or an Efflorescence, on the Skin, very like the Meafles and Scarlet Fever, but which have constantly disappeared on the Eruption of the Small - Pox; and I believe M. Gatti's Measles and Scarlet Fever, were of the same Sort, since he says, they went off sans aucun accident considerable. It would be strange indeed, if the Measles and Small-Pox should meet in the same Person, and be attended with no confiderable Accident. In the Beginning of my Practice fuch Appearances allemed me greatly, for I learned from Sydenbam, that the worst Kinds of Small-Pox appear like the Meafles, and I should sometimes have despaired of my Patients, if the Gentleman to whom this Letter is addressed, had not favoured me with the following Observation, which I received from him in a Letter, dated August 18, 1741. - " There frequently

no-

a-

gi-

ne)

n-

ar-

oft

C-

d,

in

n-

ſe.

10

1-

ly

;

-

y

t

d

e

quently is a Kind of Rash or Efflorescence on the Skin, sometimes in one Part, sometimes in another, which goes off with the Eruption, and for which I do not alter my Method." Mr. Burgefs, in his Account of Inoculation, fays, " On the fecond Day from the Attack of the Small - Pox, is often thrown out an Eruption like Fleabites, which are fometimes fo thick as to put on the Appearance of the Scarlet Fever;" but. fays this fame Author, " I know of no Instance, in so many Years since this Practice has subsisted, where any Disorder, except the Small-Pox, has been communicated by Inoculation; therefore I think it may be presumed no fuch Thing can happen, but that the Small-Pox is a Poison sui generis, and can admit of no other Mixture. I know of one Instance where the Matter was taken ignorantly by the Surgeon from a young Woman, who coming up to St. Thomas's Hospital to be salivated for the Venereal Distemper, fell ill of the Small-Pox. Three Patients were inoculated from this Matter, and had the Small - Pox in the most favourable Manner : nothing particular happened about the Wounds. They all grew up healthy; two of them are now alive, and the third died at Sea of a violent Fever many Years ago."

I will add a Case that happened in my own Practice, which may be an Illustration of the Doctrine here contended for. Some Years since, my Advice vice was asked for a young Girl that had the Itch, and within a very few Days was seized with the Small-Pox. Soon after this Seizure the Itch disappeared, and I really intagined would have been cured with the Small-Pox; but, to my great Surprize, after that Disorder was cured, the Itch broke out afresh with more Virulence than before. Suppose any one had been inoculated with Matter taken from this Person, would the Itch have been communicated with the Small-Pox? I think not—for if the Matter that caused the Itch had been mixed and blended with the variolous Matter, both Disorders would probably have been cured at the same Time.

Left the above Arguments should be insufficient to convince Mons. Gatti of his Mistake, I will add the Testimony of Dr. Kirkpatrick, whom Mons. Gatti calls the most famous Inoculator in Europe\*. The Doctor says, in his Analysis, "We proceed to consider, whether the insectious Matter is communicative of any other Disease the Person it was extracted from might have, or be liable to. For my Part, I can with great Truth aver, That an Experience of more than twenty Years on the Subject, and the utmost Resection in my Power, have convinced me, that it never did, nor can, impart any Disease but the specific one that produced

<sup>\*</sup> Le plus fameux Inoculateur dans toute l'Europe.

h,

he

if-

en

T-

ch

e.

BL

n

n

h

e

1

duced it." Dr. Kirkpatrick then adds an Observation similar to that just published of Mr. Burges; and all these Observations united, I hope will be sufficient to convince even Mons. Gatti, that no Distemper, except the Small-Pox, was ever communicated by Inoculation. However, it may be prudent to let the Matter be taken from a sound healthy Person, that has a distinct Sort of Small-Pox, as by this Means we may be certain that the Disorder will be communicated by a Mild Pus: The Practice of taking it from a virulent or confluent Sort, seems to me to be wholly unjustifiable.

Some Notice should be taken of the four following Objections, as the Author of them, Dr. De Haen, is a Physician of great Eminence.

- 1. Variolas naturales baud multum periculi babere.
  - 2. Inoculatas par premere periculum,
- 3. Tot esse qui bis variolas sustinent, ut pauci sit securitas quæ ex inoculatione colligi potest.
- 4. Eum esse numerum hominum variolis nunquam laborantium, ut sit perpetuo metus illum insicere qui iis in æternum caruisset.

It

It must appear astenishing to every one acquainted with the Nature of this Disorder, that so celebrated a Writer as Dr. De Haen should declare - " That the Natural Small-Pox is not attended with much Danger; and that the inoculated Sort is equally dangerous." In order to confute these two bold Affertions, it may not be improper to give an Account of my own Practice, during the last eight Months, in which Period I have inoculated 88 who have had the Small-Pox by Inoculation, all which recovered, and 44 in the Natural Small - Pox, of whom 25 died in the Diforder, who all had the confluent Sort, accompanied with purple Spots\*, which Sydenbam fays, are almost constantly mortis pranuncia - The Forerunners of Death. This is a Proof, that no one need be afraid to venture on Inoculation, tho' the Natural Small-Pox, at the fame Time, should be attended with the greatest Malignancy.

De Haen's third Objection is — " that fo many have the Small-Pox twice, that the Security which may be obtained by Inoculation, is of little Value." This,

\* Many of my Patients have taken large Quantities of the Peruvian Bark and Mineral Acids, without the least Benefit; for this Sort, as Sydenbam obferves, Pestem ipsam pernicie æquabat. — was as pernicious as the Plague.

This, like the two former, is not founded on Truth, and proves only how far Persons of Ingenuity may be carried by Prejudice.—For my Part, I am clearly of Opinion, that no one even had the Small-Pox a second Time, and therefore I cannot admit even of Dr. Kirkpatrick's Concession, "That one in Ten Thousand may have it twice;" for if any one ever had the Small-Pox a second Time, I can see no Reason why, it should happen only to one in Ten Thousand.

----

De Haen's fourth Objection, from the Number of those that never have the Small-Pox at all, seems to be sufficiently answered, by what has been already quoted from Riverius, Sennertus, &c. Authors that, I imagine, this ingenious Physician must be well acquainted with.

Enough having been faid in answer to Objections, which seem to be only the Cavils of a Person that is an avowed Enemy to the Practice of Inoculation, I will endeavour to consute One of another Kind, and indeed of the greatest Consequence; for it is said, and very truly, that by Inoculation, the Small-Pox is brought into Places free from it, by which the Lives of many are exposed to the satal Effects of this Disorder, who might otherwise escape it.

In answer to this I reply, that if the Small-Pox is of so infectious a Nature that very few go thro' Life without having it, except fuch as feclude themselves from all Commerce with Mankind, and are thereby rendered almost useless to Society; and if it is also true, that of grown Persons more than one in four dies, as is evident by the Accounts from the Small-Pox Hospital; and of Children, if only one in eight dies, which feems to be no unfair Representation; it follows, that Inoculation, by infecting a Town with the Small-Pox, inflead of being a public Nuisance, may really be a General Benefit, as by communicating it to Persons in their Youth, at least one-half of those who would probably die of that Disorder will be preserved; and confequently, though fome individuals may fuffer, the Community in general will reap Advantage, from the frequent Introduction of it into populous Places.

If the Small - Pox, like the Plague, could be kept from a Country or Town for Ages, we ought to guard against it by every Method that Human Prudence and Foresight can suggest; but as we know, from the Experience of at least Ten Centuries, that scarce any Country, where this Disorder has ever been introduced, can be secured from it more than sisteen, or twenty Years at most, and the sooner it comes the less Injury attends it,

then it follows, that the Inhabitants ought to be defirous of having it spread amongst them every two or three, rather than every twenty Years.

Though this Argument to me feems to be conclusive, yet if any one could convince me of the Fallacy of it, I would gladly recant, for I should be very forry to feem to recommend a Practice that may be prejudicial to Society.

d

d

In order to evince the Difficulty, if not Imposfibility, of keeping the Small-Pox from populous Places, and to demonstrate further the great Danger that attends the Natural Small-Pox, and the Benefit of Inoculation, I will insert Extracts from the Accounts that were lately sent to Dr. Huxbam by Dr. Benjamin Gale.

"The Small-Pox, by the vigilant Execution of the Laws subsisting in the several New-England Colonies, hath never generally prevailed among the Inhabitants, except in Boston, where it was epidemical in the Years 1649, 1666, 1678, 1689, 1702, 1721, 1730, 1752, and at this present Time 1764, and where the Success attending Inoculation, after much Opposition, and Endeavours used to bring the same into Disrepute, became incontestably evident.

- "In the Provinces of New-York, New-Jersey, and Pensilvania, the like Precautions have not been taken, and the Small-Pox has prevailed in those Provinces, especially in the capital Towns, once in fix or seven Years, where Inoculation hath been practised with surprising Success, to the Preservation of the Lives of many.
- "In the Year 1721, the Number of the Inhabitants of the Town of Boston were 10,567, besides those who moved out to avoid the Disease. The Decumbents were 5,989, whereof 844 died, i. e. near one in seven: At this Time, in and about Boston, 286 were inoculated, whereof 6 died, i. e. about 1 in 48: This was the Beginning of Inoculation in New-England.
- "In the Year 1752, there was an exact Account taken by Order of the Magistrates of the Town of Boston, and rendered upon Oath, in order to remove the Prejudices and Objections made against Inoculation) of all who had the Small-Pox, either in the Natural Way or by Inoculation, and the precise Number of those who died of it in either; by which Account it appears, the Number of the Inhabitants amounted to 15,734

  Those who had the Distemper in the Natural Way, amounted to 5,544; of which 514 died

  The

The whole Number inoculated, amounted to 2,113; of which 30 died.—Hitherto Mercury had not been made Use of in Inoculation at Boston.

- "In the Year 1764, at this present Writing, the Small-Pox is prevalent in the Town of Boston: By the last Accounts, 3000 had recovered from Inoculation, in the new Method, by the Use of Mercury, and five only had died, viz. Children under five Years; so that it appears, that Deaths without Inoculation, is one in seven or eight; by Inoculation without Mercury, one in 80 or 100; by Inoculation with Mercury, one in 800.
- "The Use of Mercury in the Small-Pox, was first hinted by the learned Boerhaave. This Intimation was improved, and Mercury introduced into Practice, by Physicians in the English American Colonies, about 1745.
- "Several American Physicians claim the second Glory to Boerbaave; perhaps Dr. Thomas of Virginia, and Dr. Murison of Long Island, in the Province of New-York, may justly have merited that Honour, who have successfully practised, by the Use of Mercury, perhaps before any others, either in Europe or America.

" During the late War, the Small - Pox was brought into divers Towns in this, and the other Colonies, by Return of our Soldiers for Winter-Quarters, and by Seamen employed in our Navigation to the British Islands; whereupon Application was made to the Legislature of this Colony, for Liberty to inoculate, and erect Hospitals for that Purpose, which was granted; but the Small-Pox having been thereby propagated, the Practice was prohibited on very severe Penalties, and the Inhabitants are now in a great Measure deprived of the only Method ever discovered to the World, for escaping the Hazards attending a Disease that has made fuch Havock of the Human Species, which feems to be an Invalion of the natural Rights of Mankind, and an Obstruction to their pursuing the Law of human Nature.

"The well peopling the Colonies, is a Matter of great Importance to our Mother Country; for which Reason, I cannot but think, whenever the State of public Affairs will permit the Parliament of Great-Britain to advert to the peopling and securing the Acquisitions made in America, they will judge it best effected, as much as may be, from our Colonies, and that the Law prohibiting Inoculation will be annulled, by their superintending Authority, as prejudicial to the Population of the Colonies.

" It appears from Dr. Donglas's Account of the Small-Pox in the Town of Boston, where he lived, and made critical Observations the three last Times that it was epidemical there, viz. in the Years 1721, 1730, and 1752, that the Number visited with the Small-Pox in the Natural Way was 16,047, of which 1858 died; and that in 1752, the Number of those who received the Infection by Inoculation, before Mercury was made Use of at Boston in Inoculation, were 2113 Persons, of which 30 died; granting that those who had the Disease in the Natural Way flood an equal Chance for Life with those who were inoculated, it appears, in those three Years there died 1831 in the Town of Boston only for want of Inoculation, by which Deaths, according to the longest Term of doubling the Number of Inhabitants in America, in one Century from those Periods, the Number will be diminished 29,296, which is (from the best Calculation I am able to make) a Number far superior to those who came from Europe into the News England Colonies, from their first Settlement to this present Day.

"Various Sea - Port Towns, as well as some Inland Places, have been visited with the Small-Pox, since the first Settlement of the New-England Colonies, by which many have died; perhaps, taken collectively, not more than one in four has recovered.

recovered, partly owing to removing the Sick to prevent communicating the Infection, the Want of skilful Physicians, and faithful experienced Nurfes, to attend the Sick; the Inhabitants being often obliged to make Use of some of the most abandoned Wretches, for want of suitable Persons to perform that Service.

"It is much regretted by many, that the Practice of Inoculation may not be tolerated in the New-England Colonies. As the Increase of Mankind has been more impeded by the Small-Pox, than is usually imagined, it might, perhaps, be thought Wisdom to address the Throne for Liberty to erect one Hospital in each Colony for that Purpose, that those at least who are engaged in Trade and Navigation, might obtain it, and be exempted from the Hazard of the Disease, while necessarily engaged in Business abroad.

"With regard to the Medicinal Preparation, the learned Boerhanne fays, "The specific Corrector ought to be some Medicine directly opposite to the contagious Poison: That such a Specific may some Time or other be found out, we have some Hope; and the great Use it would be of to Mankind in general, ought to encourage the diligent Part of Chymists to search for it, in some Preparations of Antimony, and Mercury."—This last Medicine may, perhaps, be justly esteemed

the most efficacious to correct the variolous Contagion, and prepare the Body to receive the Infection with Safety, that the whole *Materia Medica* is known to afford; it is therefore the Basis of the several Preparations used by the most eminent Inoculators in *America*.

The above is a Summary of Dr. Gale's historical Account of the Small-Pox as transmitted to me by Dr. Huxham; to which is added, The Method of Inoculation, as practifed in North-America; but which, not being very different from my own, as published in the former Part of this Treatise, I will not trouble my Readers with it.

Having sufficiently considered the most popular Objections against Inoculation, and, I hope, confuted them in a satisfactory Manner, I will only add a few more Instances of its fignal Advantages, that I have been savoured with by my Friends, as I am sensible the best, if not the only Recommendation of this Practice, must be taken from its Success.

Dr. Huxbam says, "He has hitherto lost no one Person committed to his Care for Inoculation, tho' he has inoculated several."—I imagine his Practice in this Way must have been very extensive, seeing he informs me, that he began to inoculate in the Year 1730. Though the Benefit of it has R 2

THILLS

been experienced so long in this County, and even under the Sanction of such an eminent Physician, yet still in many of its Districts the Prejudices against it are very great; but I hope they will in some Measure be soon removed, and in order to effect it, I beg due Credit may be given to the sollowing Testimonies, as they all have been communicated to me by Gentlemen of undoubted Honour and Veracity.

Dr. Baker, of London, fays, in a Letter I received from him lately, "That every Person in and near that Metropolis, is now entirely for it, and that it is practifed with universal good Success."

Some Time since I sent a Letter to Dr. Wall, desiring an Account of Inoculation in Worcester, and received from him the following Answer.

"Inoculation, in this Country, has been attended with very great Success, not one in many Hundreds having failed. In those who died, it has generally been found, that the Patients being very desirous of undergoing the Operation, had concealed the Particulars that they imagined might prevent the Physician, or Surgeon from undertaking it, among whom were two Women with Child.—In regard to my Method, I usually draw

## OF IN OCULATION.

off a little Blood, in order to judge of its Crasis, and if I do not find its Texture too loose, I am never fearful of undertaking the Operation; but if it should appear to be in a loose dissolved State, I prepare the Parients with the Bark and Mineral Acids.

n,

-

n

e

"Exclusive of the different Manner of communicating the Disorder, in the Natural and Artificial Infection, I have been sometimes led to think, that perhaps one Reason why the Small-Pox is milder by Inoculation, than in the Natural Way, may be because it is communicated by mild Pus: Whereas by the Natural Infection, it is received into the Lungs and Stomach by a putrid Halitus, which (as is the Case in all Diseases arising from the Stench of putrifying Substances) will contaminate the Fluids, and bring on a putrid Fever.

"An ingenious Gentleman of my Acquaintance is of Opinion, that where the Thread is to be kept long before it is used, it may be best to dry it a little before it is secured in the Phial from the external Air, as by this Means any putrifactive Fermentation in the Pus may be prevented. I have been told, that Animalcula have been obferved in the Thread when put in a little Wet without this Caution, and kept a considerable Time; but great Care ought certainly to be taken, lest by this Means the Pus should lose so much of

R 3

its Efficacy, as either not to communicate the Infection, or to produce only an incompleat variolous Eruption when the Pustules die away without coming to Maturation. I confess here, however, that though no variolous Eruption should come on, after the Inoculation, yet, if the Incisions discharge properly, the Patient may rest satisfied: For this Discharge is purely variolous; and I have known Instances of Persons inoculated with the Matter from the Incisions, who have had the Small-Pox in Consequence of it."

Mr. -- Ford, a Surgeon, near Totnes, inoculated himself, his Wife, and his two Children, in May 1742, and had fo flight an Eruption, that no one would have taken it for the Small - Pox, which made him very uneafy, as he thought he should still be liable to the Infection in the natural Way; but on my viewing the Incisions, and finding them discharge plentifully, I assured him he might be satisfied, for that he was as secure from any further Infection as if he had a great Number of Pustules. My Prognostic has been fully verified, he having acquainted me, that he has attended more than 200 Persons in Inoculation. besides several in the Natural Way, without receiving any Infection, or having the least Apprehenfion of it. He acquaints me besides, that all his inoculated Patients went through the Diforder without any dangerous Symptoms. This is one Instance,

word shade how

Instance, among many, that may be produced, that the Small-Pox, by Inoculation, even in the flightest Manner, is Security fusicient against any future Infection. visual san mon months I mai would

È

Mr. Gould, a Surgeon of Southmolton, fays, " that from the most exact Account he can procure, he finds that 293 have been inoculated in his Neighbourhood, of whom he inoculated 178, who all did well, and of the others only one died. He adds, that in the Spring of the Year 1762. the Small-Pox was fo fatal, that 9 died of the first 25 that had that Distemper, which made him afraid at first to inoculate; but he ventured after fome Time, and within fix Months inoculated 120, not one of whom had any bad Symptom, tho' the Small - Pox in the Natural Way continued to be very fatal, one out of fix having been carried off by it. Among those he inoculated, he says, one Man was turned of fixty, and another more than fifty, the first of these had a very favourable Sort. and the other, though he had been a very hard Drinker, had only three or four Puftules."

Some Years fince, when the Dorfetsbire Militia were quartered in Exeter, I mentioned to Mr. White, their Surgeon, that as the Small - Pox was then very prevalent, it might be advantageous to inoculate the common Men. The Colonel, a Gen-

tleman

tleman of great Benevolence and Humanity, approving of the Scheme, it was immediately put in Execution, and with what Success appears by the following Letter, fent me lately by Mr. White from Shaftesbury:

"The Number that I inoculated at Exeter amounted to 300; one Man, a Grenadier of Capt. Chafin's Company, loft his Life thro' the Imprudence of his Nurse, who gave him (unknown to me) forty Drops of Tinct. Theb. in the Evening. and the fame Night a large Quantity of Sheeps Dang boiled in Milk, the Confequence of which was a Delirium, and notwithstanding my utmost Care, he died the next Day, which was the fifth after the Eruption. The Number of the Pustules, which were of a very fine diffinct Sort, did not exceed thirty. One Foot, whom I inoculated, was previously infected with the Small-Pox, for the Eruption came out three Days after the Operation, and of the worst confluent kind; he lived till the eleventh Day, and I had some Hopes of him on using the Cortex and Spirit of Vitriol; for the Petechiæ, which were numerous, began to difappear, but the Disease was very heavy; and I had fome Reason to think his Attendants gave him Spiritous Liquors.

" My usual Method of Preparation, as I had no Time previous to the Inoculation, for the Colonel infisted on my filling the Hospital as soon as empty)

empty) was to bleed those that were Plethoric. give them the next Night five Grains of Calomel; and a Purge the Morning following: These were generally given twice before the Eruption .- I will mention one Case, which happened while I was at Chatham, where I had the like Success as at Exeter, with a great Number of Patients. A Serjeant had a Bubo, and other Venereal Complaints, for which he took the Sublimate, and Brandy, with the Decoction of Sarfaparilla. In the Midft of his Cure he was taken ill of the Small-Pox, had as fine a distinct kind as ever I faw, and not above 20 Pustules. I concluded him loft, on a Supposition that the Globules of the Blood were too much divided by the Sublimate for a diffinct Sort, but was agreeably deceived by his speedy Recovery."

Mr. Hodge, Surgeon, of Honiton's Clift, informs me, "That he has inoculated 300, from three Months to fifty Years of Age, and tho' feveral of them were very unpromising Subjects, of scorbutic, strumous, and leprous Habits, yet, he says, not one of them was in any Danger, which he thinks the more extraordinary, as the Small-Pox in the Natural Way at the same Time made dreadful Havock, two out of seven, and in some Families, two out of sive, having been carried off by it. Mr. Hodge adds, that he generally prepared his Patients by giving them Calomel, Æthiop. Mineral.

## 52 OF INOCULATION.

neral. Antimony, and Purgatives, and feldom omitted to bleed all above feven Years of Age."

An eminent Physician sent me, a few Days since, the following Account from Ireland.

"Although Inoculation has been a very common Practice in Dublin for many Years, yet I have no Materials from which to form even a Conjecture of the Numbers that have undergone, or the Success that has attended it. However, as I rarely hear of its being followed by any ill Event, and as the Vogue of it appears still to encrease, I believe it is as little chargeable with discouraging Consequences here as in other Countries: For, as to what we have been told from the Pulpit, of Mr. Ranby, I will only say, that I am very consident no Man that ever went before him had such good Luck in the Business: I say good Luck, because Accidents must sometimes happen, which no Skill or Caution could prevent.

"About one hundred and forty have been inoculated under my Direction, of whom one only died; but the Event was plainly not imputable to Inoculation; and I must trouble you to hear the Story.

His College of the civil and the College at the college with

103 and leis as the well disself

" A Girl, seven Years old, had such uncommon irritable Nerves, that her Parents used to call her their Hyfteric Child. She was inoculated, after the usual Preparation, in temperate Weather, from a very mild Pock; and continued perfectly well until the eighth Day afterwards. The previous Symptoms then came on regularly, and were unattended with any unufual or alarming Circumflance. The Evening before the Eruption was expected (the tenth from the Inoculation) the was calm and chearful. At this Time the was violently frightened by the fudden and horrible Screams of her Brother, who lay in the next Room, inoculated at the same Time, and of whom she was passionately fond. She fell instantaneously into a Fit of Roaring and Screeching; faid they were killing her Brother; foon loft her Senses; and flill continuing raving of her Brother until her Death. which happened on the fourth Day of the Eruption. The little Boy, who was a Year younger, of the fame Make, and, to Appearance, of the fame Habit of Body as her's, had the Disorder in the gentlest Manner imaginable.

"The fittest Period for Inoculation is evidently, I think, from two Years and a half or three Years of Age, to Puberty. I never permitted it to be practised on Infants: For, 1. They usually have the Natural Pock of as benign a Kind as the Arti-

## 54 Of INOCULATION.

Artificial. 2. While they are at the Breast, the Soreness of their Mouths (if they are beset with Pustules) is apt to disable them from Sucking. 3. Their frequent Dysenteric Fluxes make a considerable Objection to the inflicting of the Diftemper by Art. 4. From the Beginning of Dentition (which is generally between the fourth and fifth Month) till they have got their first twenty Teeth (which does not usually happen 'till they are two Years and a half or three Years old) every Practitioner knows how very fuddenly, and without any previous Notice, they are often knocked down with violent and alarming Symptoms from Teething. 5. Children, during the Period I have been speaking of, are seldom tractable. 6. It is found true, in Fact, that the artificial Infection is much more fatal to Infants than to Children from three Years upwards.

"With regard to the Preparation for Inoculation, and the Manner of performing it, the following is the Method that I always use.

"If the Patient is not in perfect Health, I bring him to it before I permit him to undergo the Operation. If there is any Acrimony in his Juices, it should be corrected by the usual Remedies, particularly Mercurials.

edited by M a calcad as

"I choose to take the Infection from a natural Pock, because the Artificial one appears to me to be less communicative of the Disorder, and more apt to disappoint us.

"The Pus ought to be taken, in my Opinion, from a young Subject, particularly if the Patient be a Child. I have feen Children inoculated from Adults, who had the Diforder in the gentlest Manner, and yet the Pock that ensued was of such a rank Nature, as to give me a Suspicion that the Practice was wrong.

"Though the variolous Matter preserves its Infection for more than a Year, yet, as all the Humours of the Body, healthy or morbid, grow putrid by being exposed to the Air, I think it obvious that fresh Pus should be preserved.

The Matter inferted should be perfectly pure and ripe. I have seen Surgeons press the Pussule until the Blood slowed; so that a Sanies was inoculated, instead of Pus. This, I have observed, is more difficult to assimilate, and the Pock that follows is more stubborn than usual.

"Although I am confident that no other Diftemper is communicable with the variolous one, yet, to content Families, I always take the Matter from a healthy Child, born of healthy Parents.

As"

"As soon as I have Notice of a mild Pock to take the Infection from, (which may be five or six Days before the perfect Maturation of it) I confine my Patient to his House, to prevent his catching the Disorder in the Natural Way; I oblige him to live temperately on plain, light Food; and restrain him from all over - heating Exercise. His Body is to be moderately loosened twice or thrice before the Operation is performed, and, on this Occasion, I would prefer gentle Mercurial Purgatives. We used, you know, to observe at St. Thomas's Hospital, (as Dr. Mead did) that all those Patients, who were taken with the Small-Pox during, or after a Mercurial Salivation, had the mildest Sort.

"If a fanguine, rigid Habit of Body, a Dispofition in the Patient to Hæmorrhages or Inslammations upon being seized with severish Complaints, or any other Circumstance indicates Blood-letting, I would choose to have it performed two or three Days before the Inoculation.

"The Patient should be inoculated in two Places, lest one should fail.

Citi rectio on tem anchieva est l'appetat à co leno anchi trevi and drive communicates at transce "I direct that the Incision should be superficial, lest the Pus should be washed out of the Wound; and that the Quantity of Matter inserted should be very small; although I must acknowledge, that I have more than once seen a large Quantity of Pus insused into a deep Wound, without the least ill Consequence.

"After the Operation has been performed the Patient should be confined to his Chamber; if he's not regular in the Discharges by Stool, his Body should be occasionally opened by an emollient Glyster; his Legs are to be bathed every Night in warm Water; and he is to wear an additional Pair of Worsted Stockings, large enough to draw half Way up the Thigh. He should, by all Means, be kept from violent Exercise; for I never saw an Instance of the Small - Pox appearing after great Fatigue of Body, that was not of a bad Kind. His Allowance of Flesh Meat should be gently lessened a Day or two before the expected previous Fever, and light Puddings, Broth, or Gruel, substituted in the Place of what is cut off.

"After the Invalion of the previous Symptoms, the Disorder, you know, is to be treated in the same Manner as the Natural one; but more Purging seems to be required after the Pock is dry, to prevent the Consequence of Boils."

The following Account I received from a Person of great Rank and Fortune in this Country.

" A gentleman engaged in the public Service of his Country, having often lamented that his Attendance thereon was prevented by an Apprehension of the Small-Pox, resolved on Inoculation, tho' aged, and of a corpulent habit; and having happily recovered of that too commonly fatal Diftemper, in humble and grateful Acknowledgment of fo great a Bleffing, caused an Intimation to be given, that he would defray the Expence of all within his Parish who were inclined, and should be deemed fit, to be inoculated; in consequence of which eighty-fix underwent the Operation in the Parish of King's-Nympton, every one of whom recovered of the Small-Pox, except one Child that was feized with a Fever the Day after the Operation was performed. It is remarkable, that among those inoculated Persons there were several aged, and some who were of the most scrofulous Constitutions."

In Exeter, according to the best Calculations I can make, since the Year 1741 there have been about 700 Persons inoculated, of whom only two died; one of these I have already given an Account of and the other was not my Patient.

It feems needless to mention any other Instances, as Inoculation is now practifed with the greatest Success in almost every Kingdom in Europe, and is strongly recommended by the whole College of Physicians in London, as appears by the following Judgment, which they ordered to be entered fome Time fince in their Annals-" Quoniam Collegio nunciatum fuit, falsos de variolarum insitivartm in Anglia successu, et existimatione apud exteras gentes nuper exiisse rumores, eidem collegio sententiam suam de rebus hisce ad huc modum declarare placuit, (viz.) Argumenta, quæ contra banc variolas inserendi consuetudinem in principio afferebantur, experientiam refelisse, eamque boc tempore majori in honore apud anglos haberi, magisque quam unquam antea inter eos nunc invalescere, atque bumano generi valdi salutarem esse se existimare." --- " The College having been informed, that false Reports concerning the Success of Inoculation in England have been published in Foreign Countries, think proper to declare their Sentiments in the following Manner, viz. " That " the Arguments, which in the Beginning were " urged against it, had been refuted by Experience; " That it is now held in greater Esteem by the En-" glish than ever; that the Practice of it increases " greatly among them; and that the College think " to be highly salutary to the Human Race."

Exeter, JOHN ANDREW.
May 17, 1765.

ONTHE PROPERTY OF A 2 域。其特別的國際問題,但可以 er mig bei bei gem bar ben es bis in begenere school and the state of the state of the post of party and Carlos de la companya del companya del companya de la companya de William to a limit the willing A THE SHOP IN THE SHAPE OF THE Hilliam Miles Assistant and March and And the facility of the same of the same in the same of the same o William Feld and the share the state of the the transfer Mary is not been a series of the same 



## POSTSCRIPT.



O O N after the foregoing Treatife was finished, an unfortunate Event happened at some Distance from Exeter, of which it would be unpardonable in me not to give a true Account.

On the 21st of last Month I was desired to attend the Inoculation of two Children, the Operator having never seen any Thing of the Practice. One of them had been under my Care, a Fortnight or three Weeks, but the other I never saw till the Time of the Operation.

Being

Being present when they were inoculated, I said I would come and see them during the Eruptive Fever, (as is usual) but, to my great Surprize, my Offer was rejected, and therefore I no longer considered them as my Patients. However, I was called to them the 2d Day of this Month, (which was the thirteenth after the Operation) when I found both ill of the consuent Small-Pox. The Eruption being at that Time completed, it was too late to prescribe any of the Evacuations, which are generally ordered on the Seizure of the Small-Pox, if it proves violent.

From the 2d of June, I had no particular Account of them, till the 6th, when I visited them, with my Friend and Colleague Dr. Glass, which was the only Visit we made the youngest, \* for he died in the Evening of the following Day; but we were desired to attend the eldest again, on the 8th, at which Time there appeared Hopes of his Recovery; and favourable Accounts of him were sent to us till the 14th, when we were acquainted that he had brought off six Worms the Day before, and that he was in the last Extremity.

As

\* A Sifter of this Child was inoculated under my Directions, at the same Time, but in a different Part of the Country, and had the Small-Pox in a very favourable Manner.

As neither Dr. Glass, nor myself, ever saw such a malignant Sort of Small-Pox from Inoculation, among near a thousand that have been inoculated under the Directions of one, or other, of us, as it resembled exactly the virulent Kind, which has greatly prevailed during the last fix Months; and as these Children had been almost surrounded by the Small-Pox, it may not, perhaps, be unjust to suppose that the natural Infection had reached them. I wish their afflicted Parents could be of this Opinion, for then they might be fatisfied, that the same Fate would probably have attended them, if they had not been inoculated. On the Supposition, however, that the Deaths of these Children may be imputed to Inoculation, and that both of them ought to be deemed my Patients, (which, for the Reasons already mentioned, I think would be unfair) yet three only have died of about 600 that have been inoculated under my Direction. How disproportionate this to the Loss by the Natural Small-Pox?

It is reckoned by some ingenious Calculators, that one in 12000 dies every Day; and as the Term allowed for Inoculation is reckoned to be 21 Days, the inoculated Patient's general Chance of dying, during that Period, would be nearly as one to 571, abstracted from the particular Hazard attending

attending that Practice, which ought to be admitted, to account for the Deaths of some Persons under Inoculation; and probably Worms might have contributed to the Death of the Child last mentioned, for besides the six brought off the Day before his Decease, he voided several during the Course of his Disorder.

Notwithstanding what is advanced in Page 42, and 43, I beg it may be understood, that I do not intend to recommend the Practice of Inoculating in Towns that are free from the Small-Pox; and in fuch as are infected, it would be very hazardous; for which Reasons it ought never to be performed, except in remote Places, appropriated to that Purpose.

Whatever Inelegancies, or Inacuracies may occur in this Treatife, the Reader, it is hoped, will be candid enough to excuse them, especially when he considers, that the Writer had no Leisure to polish, or correct his Style, being only attentive to the Facts, and which he committed to the Press amidst the Multiplicity of indispensable Avocatitions, without the least Ambition of becoming an Author; this fole Motive being to obviate the Prejudices which have obstructed the Practice of Inoculation, in the City and Neighbourhood of Exeter-a Practice which he is fully convinced, (notwith(notwithstanding some fatal Accidents that must necessarily attend it) will be beneficial to Mankind, in proportion as it is extended.

June 17, 1765.

An Account of the Inoculation of the Duchess

[Inoculation for the Small-Pox having met with great Opposition in France, any little Miscarriage that happens from Ignorance or Inexperience, is misrepresented by its Enemies, as a just Reason for prohibiting the Practice. The Case of the Duchess of Boufflers gave the Opposers great Advantage; It was therefore necessary for her Physician, one of the most eminent in France, to state her Grace's Case, as it really happened; by which he is forced to confess his own Mistake, in order to undeceive the Public. The Affair has made a great Noise abroad, but has not been generally underficed here, and therefore, the following authentic Account is inserted.]

And about the levente or ciclica Days

Translation of Mons. Gatti's own Letter to a Friend on the Subject.

SIR,

T is but too true, that the Duchess of Bouf-I flers, whom I inoculated, about two Years and a half ago, is just recovered from the Natural Small Pox, wich proved to be a diffinet and favourable Sort. This event having greatly excited the Attention of the Public, I thought it my Duty to give the World a particular Account of the Duchess's Inoculation; and she has, upon my Application, been pleafed to recollect all the Circumstances, and to favour me with the following Certification figned with her own Name.

"On the 12th of March 1763, I was inoculated for the Small-Pox; and about four or five Days afterwards, a Redness appeared round the Orifice. which Monf. Gatti called an Inflammation, and affured me was a Sign that the Small-Pox had taken Effect. These were the very Terms he used. This Redness or Inflammation increased every Day. and about the feventh or eighth Day, the Wound began to suppurate. There appeared also about the Wound fix small Risings, or Pimples, which fuccessively suppurated, and disappeared the next Day.

Day. Monf. Gatti, upon these Appearances again affured me, that the Small-Pox had taken Effect. In the Afternoon of the eleventh or twelfth Day of my Inoculation, I felt a general Uneafiness and Emotion; a Pain in my Head, and my Back, and about my Heart; in Consequence of which, I went to Bed sooner than ordinary; I slept well, however, and rose without any Disorder in the Morning. These Symptoms Mons. Gatti affured me were the Forerunners of the Eruption. The next Day a pretty large Rifing, or Pimple, appeared in my Forehead; which came to a Head, turned white, and then dried away, leaving a Mark which continued many Days.

The Wound in my Arm continued to suppurate seven or eight Days, and Monf. Gatti now affured me that I had nothing to fear from the Small-Pox; and upon this Assurance I relied without the least Doubt, and continued in perfect Confidence of my Security till the Natural Small-Pox appeared. I continued very well during the whole Time of my Inoculation, except one Day, as mentioned above, and I went out every Day.

MONMORENCY, D. de Boufflers."

Calson Device dela Tale to the off the

Contraction of the State of the

not tee stations, and I store is not tell eachies of

00

It is very true that I affured the Duchess, she had nothing to fear from the Small-Pox; but the Event has proved that I was mistaken.

I confidered the Aceidents or Symptoms which followed the Incision, as certain Signs that the Inoculation had taken Effect: As the characteristic Symptoms of the inoculated Small-Pox did not follow, I mean a Fever succeeded by a Suppuration of the Wound, a Suppuration different from that which might take Place before the Fever, I thought that the Action of the variolous Virus could produce no other Effect in the Patient, than what appeared round the Incision, and that she was therefore secure from the Small-Pox; and I was confirmed in this Opinion, by her continuing exposed to the Contagion with her Daughter and another Lady who were inoculated at the same Time, and had the Small-Pox with the usual Symptoms.

I was persuaded, according to the general Opinion, that when Signs of the Small-Pox having been taken, appear round the Incision, that Disease will certainly follow, supposing the Patient to be susceptible of it; and that if the Disease does not then follow, the Patient is not susceptible of it. The Case of the Duchess proves that this Opinion

nion is false, whether the Signs of the Small-Pox having been taken or equivocal, or whether the Virus may act upon that particular Part without spreading to the rest of the Body. I make no Account of the Pimple which appeared upon her Forehead; because it had neither the Form, or the Course of a variolous Pustule; and the Duchess herself, who has now good Reason to be well acquainted with variolous Pustules, is of the same Opinion.

As the same Thing that has happened to the Duchess has happened also to many others where Inoculation is frequent, and as it may and must happen again, it is of great Importance to prevent the Mistakes and calm the Anxieties that it may produce: And for this Purpose I have always thought, that every Patient who has received the real Small-Pox by Inoculation, ought to have a Certificate of it given them by their Physician.—
I have given many such Certificates, and I should be glad if every Person whom I have inoculated would apply to me for one.

T 3 G. M.

es in a kithin simble as was



### TO THE

PRACTICE of INOCULATION.

ANDREW, Dr. John, the Method practifed by him, in Inoculation, p. 6

— Mercury, as used by him in preparing the Subject, 10

— the Diet recommended during the Progress of the Disorder, 11

— his Manner of conveying the variolous Matter, 12

— particular Treatment of the Patient, ib. 20

— Diet prescribed on the Recovery, 13

— the Time the Small-Pox is usually observed

ferved to turn, when communicated by
Inoculation, 13
what is necessary to be done, in Case
of Reftlessness, 14 — the Age most approved of, for the Ope-
ration, - 21
his Account of two unsuccessful Sub-
jects, 61
- Numbers inoculated by him, 63
- the Places most to be approved of for
the Operation, 64
Antimony and Mercury, the most promising
of the Materia Medica, to afford a Spe- cific for the Preparation for Inoculation,
Ambassador's, the French, his Observation
on the Practice of Inoculation in Eng-
8rt NDR EU. Dr. John the brait bed
and the statement of the problem in the part of the
B. Billettell
to a coject,  10  10  10  10  10  10
Baker, of London, Dr. his Account of the
Progress of Inoculation, 46
Bear's Foot, it's Efficacy in discharging
Worms from Children,
the Species which should be carefully
chose for the Purpose, (Note)
bayran — its
[18] [18] [18] [18] [18] [18] [18] [18]

- its good Effects with Calomel, in prepar-
ing four Children for Inoculation, 16
Boerhaave, Dr. his Opinion of Inoculation,
(Note)
- First hinted the Use of Mercury, 41
- His Opinion in regard to the Medicine
most effectual for a Preparative to Inocu-
lation, 44.
Boston, (in New-England) Progress of the
Small-Pox there,
- Inoculation first introduced there, 40
- Dr. Douglass's Calculation, when the
Small-Pox was epidemical there, 43
Boufflers, Duchess of, a particular Account
of the Progress of the Small-Pox, when
she was inoculated by Dr. Gatti, 66

#### C

Calomel, and a particular Species of Bear's
Foot, their good Effects in discharging
Worms from Children, preparative to
their being inoculated,
16
Clarke, Dr. his Letter directing one of his
Servants to be inoculated; with an Account of the Progress of the Disorder, 2
College

College of Physicians, their	Opinion of the
Practice of Inoculation,	.59
Condamine's Calculation of I	Lives, 28
Calculation, another,	63
and the little part of the process.	

#### D

De Haen, Dr. his Objections to the F	ractice
of Inoculation refuted,	35
Divines, the most eminent, their Opi	nion of
the Practice of Inoculation,	22
Douglass, Dr. his Observations on the	ie Pro-
gress of the Small-Pox, when epic	lemical
in Boston,	43

#### E.

Exeter	City,	Number	of	People	inoculated
e th	ere sinc	ce 1741,		ag is br	a deared 38
- an	Accou	int of a r	ema	rkable !	Miscarriage
- in	the Pi	actice the	re,	STATE OF THE	61

Charles I have med that a sufficient de deep at that

E Carron 2 of January 50 to 10 1000

F.

of a line going on the bodies.

Ford, of Totness, Surgeon, an Account of his inoculating himself, his Wife and Children, 48

Ğ.

Gatti, Dr. his opinion in regard to the Time most proper for Inoculation, 19

— his Account of the Case of the Duchels of Bousters, when inoculated, 66
Gloucester, Bishop of, his Opinion of Inoculation, 22
Gould, of Southampton, Surgeon, Account of his Success in Inoculation, 49

H.

Heberden, Dr. his Opinion of Inoculation,

30

Hodge, of Honiton's Clift, Surgeon, an Account of his Success in Inoculation there,

with

with his Method of preparing his Subjects, 51

Huxam, Dr. his Strictures in regard to the
Use of Mercury in Inoculation, 19

his Account of the Success of Inoculation
in America, 21-45

#### ī.

James, Dr. his Powder useful in preparing
the Subject for Inoculation,
Inoculation, when first discovered and intro-
duced into England, fee the Preface.
- the Case of an unfortune Subject, 14
- the Age most to be approved of for the
Operation, 18—21
- Opinion of different Physicians on, (note)
the state of the s
- fome of the most popular Objections to
the Practice obviated, 21
- its Advantage clearly shewn by Calculati-
on, 1041 10 1011 1 23
- Opinion of Foreigners in regard to the
Practice, 26
- on the Appearance of a Rash or Efflores-
cence,

cence,	different	<b>Opinions</b>	about	them,
the soul for	letognist.	ali est ch	distribution of the second	33
- a rema	rkable Inst	ance of th	e Safety	of the
Practic	e,	i every	计多层部	26
- Calcula	tions, she	wing the	Value o	of the
Practic	e to the H	luman Spe	cies,	39
- introdu	ced into N	lew Engla	nd, (at B	ofton)
a remai	rkable Acc	ount of i	s good I	Effects
in the e	early and p	refent Pra	dice,	40
- Accoun	t of, the	great Im	provem	ent of
the Pra	ctice by	the Introd	uction .	of the
Use of	Mercury,			41
- Advant	age to th	e Practice	of, by	a just
Compai	rison of the	e Number	who hav	ve had
the Sm	all Pox in	the Natu	ral Way	, &c.
- 524 323116	TOPPOST OF	Doctoria	dirick,	43
Ireland, an	eminent	Physician'	s Accou	int of
the Pra	ctice of In	oculation	in,	52
- his Cafe	e of a Sub	ject that r	niscarrie	d, 53
- the Tin	ne and Ag	e approve	d of by	him,
with fo	me Obser	vations no	eceffary	to be
attended	d to,			ib.
- his Mar	ner of pr	eparing Su	ibjects,	54
- the Var	riolous Ma	itter to b	e taken	from
the Nat	ural Pock,	as being	most con	nmu-
nicative	, and fron	a young	Subject	55
1	Calengal d	Julius &	18 4	State
THE SHARE		The second		

_	State the Pock should be in when	n the Pus
	is extracted, for the Purpofes of	
A	tion, it is to the state of the	55
_	Subject how to be prepared,	56
-	the Incision should be made super	erficially,
	and why,	57
-	Confinement necessary after the on, with regard had to the Habit	
100a.	Treatment of the Subject, on t	
	off of the Diforder,	57
215		A 10

### K.

Kirkpatrick, Doctor, his Account of	f the Suc-
cefs of Inoculation,	20
- his Opinion in regard to comm	unicating
it, it,	34

# M. .

Mackinzie, Dr. his Opinion in regard	to the
best Time for Inoculation,	19
Mercury, its Use in Inoculation,	7
- Dr. Wall's Strictures on its Use,	ib.
	Dr.

#### IN-DEX

4 2.5	
-	Dr. Huxam, on the same, with the great
++1	Advantage arising from it,
-	the Use of it very falutary in Inoculation,
	10-15-16-41
-	the Use of it first introduced by Dr. Boer-
	haave, Dr. Thomas of Virginia, and Dr.

#### N.

Murison of Long Island,

Norwich, Bishop of, his Opinion of Inoculation, 22 New-England, obstinacy of some of the Colonies there against the Practice of Inoculation, 44

0.

Objections of Dr. De Haen to the Practice of Inoculation refuted,

U 2

Powder

### INDEX,

No.		340		ing.	<b>P</b> (\$0.5)	(8年)	(1)	El a	100	oko i	議員		to the
e la	, de		1.50		, 12	P		Tie		tgia.	ar.	Male :	
op.		0	trio.									confirm	

Powder, Dr. James's, against Fev	ers, its
Use in preparing the Subject for	
tion,	11
Pus, the Kind best to be chosen, and	how to
be preserved for Use,	47

# able of the continue of the Carlotte of the Market of

parties and the second

Scheuzer, Dr. his Account of the Success	of
Inoculation,	20
Small-Pox in the Natural Way, how dange	er-
ous,	24
- of the almost Impossibility of escaping	it, b.
- Difficulties attending the Recovery,	30
- Calculations chiefly shewing how destru	c-
tive it is to the Human Species,	39
Squire, Dr. Bishop of St. David's, his Or	oi-
nion in regard to Inoculation,	28

tabulat in I for a file

Young

- 8. Care must be taken, during the great Heats, that the Soldier on Duty should remain exposed to the Heat of the Sun as little as possible, and that he avoid sleeping there. The Cuirassiers, especially, when once their Cuirass is thoroughly heated, are those who suffer most by the Sun.
- 9. Neatness cannot be too much insisted on. Let Soldiers frequently wash their Hands, their Face, and their Feet; and if the Season permits, let them bathe as much as possible in running Water.
- no. Great Care ought to be taken to lodge many men together in a small Space; and if it cannot be avoided, let the Air be at least renewed as often as it can, whether those who lodge together are in Health or Sickness; for from hence arise the most dangerous, and even the contagious distempers.
- made of good and pure Flower; for it is certain that very dangerous Distempers are occasioned by musty or spoiled Bread.

b

CON-

minastral range (1) and the first of the fir

ngšila († 1862) 19. standards proportion in Papada († 1862) 19. standards proportion († 1862) 18. standards proportion († 1862)



# CONTENTS.

# LILLILLIA

INTRODUCTION	page 1
Of Coughs	2
- a Sore Throat	3
- the Pleurify	3 8
- a Peripneumony	17
- the Rheumatism, and Rheumatic	Pains
4.	30
- Intermitting Fevers	35
- Spring Intermittents	38
- Autumnal Intermittents	43
- Quartan Fevers	48
- the Jaundice	49
- the Dropfy	51
- Vomiting	. 58
- the Colera Morbus	ib.
- a Diarrhæa	61
- the Dyfentery	63
	Of

# CONTENTS

Of an Inflammation of the Intestine	s 68
- the Phrenzy	72
- the Hæmorrhage of the Nose	76
- a continued Fever	0 19
— the Scurvy	86
- the Gangrene	92
- the Lues Venerea	93
- the Itch	99
— Worms	102
RECIPES, a Collection of,	properly
numbered, so as with Ease to b	CONTRACTOR OF THE PERSON OF TH
to when mentioned in the Courf	e of the
Work.	105



special granting persons

Continues of principle

en salatan kerenangan -- Georga Keras

Scibara ( ont --

the Dieply

miting -

- the Colon Ma - a Diardien - the Dyenery

00

33

82 54

.

(H)

7.0

88

63

Young, Dr. his Opinion in regard to the most proper Age for Inoculation, 19

#### W.

Wall, Dr. of Worcester, his Sentiments in
regard to the Use of Inoculation,
- his Account of the favourable Advances
in the Practice of Inoculation, 46
White, Surgeon of the Dorsetshire Militia,
a remarkable Instance (in his Practice) of
the happy Effects of Inoculation, with the
Preparation he was once constrained to
put in Practice, 49
Worms, Danger of them to an inoculated
Subject, 10
- Bear's Foot, its Efficacy in expelling
them, with the Species to be chosen, ib.
- fuspected of occasioning a Miscarriage in
a Patient, 64

Well Dr. of Marcelle. his Stationed in recommendation of the culture on the contract of the culture on the culture of the cult



## RECIPES

Referred to in the foregoing

### TREATISE.



I.

TAKE, of the Species for the Pectoral Decoction Three Ounces; boil it in a sufficient Quantity of Water for Half an Hour, to strain Three Pounds.

2.

Take, of the Mass for Pil. Storax, Eight Grains. Make two Pills for a Dose,

Allowed in the Prediction of

Take,

3

Take, of the Species for the Emollient Decoction Six Ounces; boil it in sufficient Water to the Thickness of a Cataplasm\*, adding toward the End One Ounce of Mustard Seed bruised, for a Poultice.

4.

Take, Flowers of Elder One Ounce; let it just boil up in a sufficient Quantity of Water in a close Vessel; then digest warm for Half an Hour; strain off Two Pints; to which add an Ounce and Half of Rob of Elder, and Forty Grains pure Nitre; Mix them together.

5.

Take, Flowers of Elder, and red Roses, of each Half an Ounce, pure Nitre One Drachm; mix them together. Insuse a Pugil+ of this in warm Water for Tea.

б,

Take, Leaves of Senna Six Drachms, Water Scurvy-grass Two Drachms, Agaric One Drachm, and

<sup>\*</sup> A Poultice.

<sup>+</sup> The Eighth Part of an Handful.



### TO THE

DISEASES OF THE ARMY.

### CEDECECECECECECEC.

A.

AUTUMNAL FEVERS, how to be treated, 43

B.

Bleeding at the Nose, Directions for treating,
76
Bread, a Precaution necessary to be attended
in regard to, (Preface)

Buboes, when they grow hard, how to be
treated,

M Camp

C

Camp, the best Situation for one, (Preface)
- frequent Movement of which contributes
to the Preservation of the Troops, xii Cautions necessary to be attended to, in order
to preserve Health amongst the Troops, ib.
Colera Morbus, or a fudden Evacuation both upwards and downwards, 58
- Directions for what is immediately to be
done, 59  Directions to be attended to on the first
Abatement of the Disorder, and the Food necessary to be taken,
Coughs, a Variety of them, the Manner they
are to be treated, with the Regimen ne- cessary in the Complaint, 20

#### D.

Defign of the Publication,	(Preface) viii
Diarrhæa, what, and how d	listinguished from
the Dyfentery,	61
- when of Service to the	Preservation of
Health, and otherwise,	ib.
AND CONTRACTOR OF THE PARTY OF	- when

- when dangerous, a Prescription for, 62
Dispensatory, 10
Dropfy, what, and how occasioned, with it
particular Appearance,
- how to be treated, and the Regimen ne
ceffary,
- when Tapping is necessary, and how to be
performed to have Effect,
- how to diffinguish this Disorder, whether
occasioned by Wind, or a Collection o
Water,
- when from Wind what is necessary to be
done,
- when the Fluid Matter is collected in the
Breaft, Signs by which it may be disco-
vered, with Directions for what is neces-
fary to be done, ib.
Dysentery, how to be distinguished, and how
occasioned, 63
- necessary Instructions to be attended to
when the Disorder prevails, 64
- how to be treated,
- when evacuating Medicines are to be dif-
continued, 66
- when frequent Inclinations to flool with-
out any thing passing, with Pains in the
lower Belly, a Prescription for, 67
En application of the second

### I N D E X

### E.

Evacuations both upwards and downwards, for Direction about, see Colera Morbus, 58

#### F.

现得来的原理的现在分词 化多种甲基酚甲基二甲甲基二甲甲基酚酚甲基二甲基酚甲基甲基酚甲基酚甲基酚甲基酚甲基酚甲基酚甲基酚甲基酚甲基酚甲基酚甲基酚甲基
Fevers, Intermitting, the different Species of,
and the apparent Signs they are attended
with,
- the Spring Intermittent, manner of treat-
ing it,
- Directions in regard to the Food to be
taken, 39
- different Prescriptions necessary to be at-
tended to, according to the Circumstances
of the Patient, 40
Fevers, Autumnal Intermittents, their diffe-
rent Appearances, 43
- Prescriptions necessary to be followed,
distinguishing the Cases, 44
- the Food necessary during the Disorder, 45
- when the Fever is obstinate, a Prescripti-
on for,
Fevers, Quartan, its Appearance and how to
be treated, 48
Fevers,

Fevers, continued, its Symptoms, 79.
- what is necessary to be done in this Difor-
der, 80.
- the Signs which indicate an Abatement in
the Diforder, 81
Fevers, the continued, or Hospital Fever,
how generated, 81-82
- favourable, and unfavourable Symptoms,
83
- Prescriptions necessary to be followed, 84
Forces when in encampment, precautions ne-
ceffary to be attended to, with an Account
of the Disorders they are incident to,
Control of the second s

### G.

	fe in preferving People in Health and Vigour, (Pref.)	
Gangrene, necoffary	Prescriptions to prevent	4
or remove it,	92	2
Gonorrhæa, how to	o be treated, 27	7

#### H

Hæmorrhage,	or Bleeding of	the	Nofe, dif-
ferent Ten	dencies from,		76.
	*M 3		- Signa

- Signs which declare it falutary, and	
otherwife,	ib.
- different Method of removing it,	77
1.	
Jaundice, its Appearance and how to be	reat-
ed,	49
Inflammation of the Intestines, the Symp	toms
by which it is discovered,	68
- the Prescription for,	69
- when a Gangrene is apprehended,	70
- what is necessary to be done on the Fo	rma-
tion of an Abscess,	ib.
Itch, Signs by which it may be distingui	fhed,
	99
- when of the inveterate kind,	ib.
- Directions for treating it,	100
- when anointing the Parts, a necessary	Pre-
caution to be attended to,	ib.
L.	
Lodging of Soldiers, a Precaution nec	effary
to be attended to, (Preface)	xiii
Lues Venerea, or Venereal Disorder,	93
- the different Complaints arising from	The state of the s
Disorder,	ib.
	ymp-
	W. 1289

- Symptoms by which the Diforder is difco-	
vered, 94	
- a safe and sure Method of treating it, ib.	æ
- Signs by which it may be discovered, that	
the Prescription has taken Effect, of	
- Regimen necessary to be followed, ib.	200
- should a Salivation come on, a Caution to	800
be attended to,	
- Signs by which a Salivation may be	650
known, ib	
- what is necessary to be done on the Ap-	
pearance of a Gonorrhæa, ib	962
- on the Suppression of the Gonorrhaa	237
should the Testicles become swelled, Di	wall
rections for,	
- Directions to be followed, when the vene	
real Buboes grow hard, which the vene	-80

### P.

Per	ipneumony, or an Inflammation on the
	Lungs,
_	the Symptoms that usually attend it in an
	unfavourable State, ib.
-	Bleeding, when necessary, and what should
	be the Consequence to shew a good Effect,
	18
	- Direc-

# I'N D'E X

- Directions for a further Treatment	of the
Patient, with the Diet he should be	kept
to,	19
- when an Abscess may be apprehended	ed, 22
- Directions for bringing it forward an	d dif-
charging the Pus,	25
- when an Abscess is formed at the I	Extre-
mities, Directions how to treat it,	28
- when the Complaint terminates in a	fchir-
rous hardness, the Consequence,	ib.
Pleurify, what, and the different Sym	ptoms
that attend it,	8
— a false one, how known,	ib.
- Bleeding, when necessary, and what	t is to
be observed when performing,	9
- Fomentation, or a Plaister of Lauda	
when necessary,	ib.
- when a fecond (or oftener) Bleeding	
ceffary,	ib.
- Regimen necessary to be observed,	10
when Respiration grows easy, Direct	tions
for,	II
- when Blisters may become useful,	ib.
- the different Symptoms that gives I	Iopes
of Recovery,	1.2
- Signs by which it may be known to	
a Tendency to a Consumption, with	
rec	dions.

rections, then necessary to be attende	d to,
Property and the second	14
Phrenzy, or a continued Delirium, atte	ended
with an acute Fever,	72
- how occasioned, and the Symptoms	with
which it is usually attended,	ib.
- Directions for the Treatment of the	Pati-
ent,	73-
- the Regimen necessary to be followed	d, 74
	the second

### R.

Rains, Time of, how the Tents in a Can	10
	kii
Recipes, or the Dispensatory referred to	in
	05
Rheumatifm, and Rheumatic Pains,	30
- the different Stages of the Disorder, a	ba
how contracted,	ib.
- Directions for its Treatment,	32
- Appearances that are favourable,	33
when the Complaint terminates in a fix	ed
Pain, how to be treated,	ib.
- what is necessary to be attended to after	a
Recovery,	34
가다는 가수요 있다는 사람들의 회의 회의 회의 회의 회의 회의 기계를 가는 사람들이 되었다. 그 사람들은 사람들은 사람들이 되었다면 하는 것이 되었다면 하는데 그 사람들이 되었다.	he
Parts, how they are to be treated,	ib.

Salivation,

S.

Salivation, what, and the Appearances by
which it may be discovered, 97
Scurvy, the Symptoms by which it is disco-
vered, 86
- how contracted, and to be treated, 87
- Regimen necessary to be attended to, 88
- Symptoms to be attended to, at the going
off of the Complaint, 90
Soldiers newly raised, how they should be
treated, (Preface) ix
- a Caution to them when heated on a
March or with Work, xii
Stools, when frequent and liquid, fee Diarr-
hæa, 6r
Deposition of the form of the property of
$\mathbf{r}$
while his experience will be able to the wife of
Throat, fore, various Kinds of, with the
most immediate Remedies necessary, 3
- when the Tonfils grow red, and the
Glands are attacked, manner of treating;
- 4
- when Bleeding is necessary, ib.
the Regimen to be taken in this Difor-
der

- a Suppuration, when brought on, how to be managed,
  when an Abscess is formed, how to be treated,
  when Deglution is stopped, the proper Glysters to be used,
- a Species of, inclinable to a Mortification, how to be treated, ib.

#### V.

Venereal Disorder, the Treatment of it in its different Stages, see Lues Venerea, 93 Vomitings, how to be removed, with some Directions necessary, 58

#### W.

Water, Directions for sweetening, when foul,
(Pref.)

Wind, a free Passage to which should be attended to in all Encampments,

Worms, Signs by which they are discovered,

102

how to be discharged,



Market 12 Walter constitution of the second All as weath the article in 120 About 100 feeting British of the American State of the State o At James of the same to the sail on the fell · 通過過過過程上於 1700年1月 · 100年1 eith to respond the firm that the second Se Wardell or the way of the fifth and Article State Company of the State of the St AND A SECTION OF THE PARTY OF T The state of the s Signal Contract Charles to the Control of the Contro to set that saile is no officed a well ary the state of t Weens, Standard with they are a transaction.